

BEST AVAILABLE COPY

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C12N 15/12, C07K 14/475	A2	(11) International Publication Number: <b>WO 00/18914</b> (43) International Publication Date: 6 April 2000 (06.04.00)
<p>(21) International Application Number: PCT/US99/21647</p> <p>(22) International Filing Date: 17 September 1999 (17.09.99)</p> <p>(30) Priority Data: 09/161,241 25 September 1998 (25.09.98) US</p> <p>(71) Applicant: AMGEN INC. [US/US]; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).</p> <p>(72) Inventors: BASS, Michael, Brian; 2208 Rustic Park Court, Thousand Oaks, CA 91362 (US). SULLIVAN, John, Kevin; 1085 Rotella Street, Newbury Park, CA 91320 (US). THEILL, Lars, Eyde; 1874 Calle Borrego, Thousand Oaks, CA 91360 (US). WANG, Daguang; 1800 Via Petirrojo #K, Thousand Oaks, CA 91320 (US).</p> <p>(74) Agents: ODRE, Steven, M. et al.; Amgen, Inc., One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>	
<p>(54) Title: NOVEL <i>DKR</i> POLYPEPTIDES</p> <pre> 1  MQRIGATILLC LLLAAAVPTA PAPAPTATSA PVKPGPALSY PQEEATLNEM 51  FREVEELMED TQHKLRSAVE EMEAEEAAAK ASSEVNLANL PPSYHNETNT 101 DTKVGNNTIH VHREIHKITN NQTGQMFSE TVITSVGDEE GRRSHECIID 151 EDCGPSMYCQ FASFOYTCQP CRGQRMCTR DSECCGDQLC VWGHCTKMAT 201 RGSNGTICDN QRDCQPGGCC AFQRGLLFPV CTPLPVEGEL CHDPASRLLD 251 LITWELEPDG ALDRPCASG LLCQPHSHSL VYVCKPTFVG SRDQDGEILL 301 PREVPDEYEV GSFMEEVROE LEDLERSITE EMALGEPAAA AAALLGGEEI 351  * </pre> <p>(57) Abstract</p> <p>Disclosed are nucleic acid molecules encoding novel <i>DKR</i> polypeptides. Also disclosed are methods of preparing the nucleic acid molecules and polypeptides, and methods of using these molecules.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## NOVEL DKR POLYPEPTIDES

5 Field of the Invention

This invention relates generally to novel genes encoding proteins that have use as anti-cancer therapeutics.

10

## BACKGROUND

Related Art

One of the hallmarks of cells that have become cancerous is the change in the gene expression pattern in those cells as compared to normal, non-cancerous cells. An intricate series of cell signaling events leads to this so called "differential gene expression", resulting in conversion of a normal cell to a cancer cell (also known as "oncogenesis" or "cell transformation"). A number of cell signaling pathways have been implicated in the process of cell transformation, such as, for example, the cadherin pathway, the *delta/jagged* pathway, the *hedgehog/sonic* pathway, the *hedgehog* pathway, , and the *wnt/wingless* pathway (Hunter, *Cell*, 88:333-346 [1997]; Currie, *J. Mol. Med.*, 76:421-433 [1998]; Peifer, *Science*, 275:1752-1753 [1997]). Interestingly, these same pathways are involved in cell morphogenesis, or cell differentiation, during embryo development (Hunter, *supra*; Cadigan et al., *Genes and Develop.*, 11:3286-3305 [1997]).

The *wnt* genes encode glycoproteins that are secreted from the cell. These glycoproteins are found in both vertebrate and invertebrate organisms. Currently, there are at least 20 *wnt* family members,

- 2 -

and these members are believed to function variously in control of growth and in tissue differentiation. Recently, discovery of a novel gene was identified in *Xenopus* and mouse and has been termed *dickkopf-1* ("*dkk-1*"). This gene is purportedly a potent antagonist of *wnt-8* signaling (Glinka et al., *Nature*, 391:357-362 [1998]). Interestingly, this gene is also purportedly involved in morphogenesis in the developing embryo (Glinka et al., *supra*). This gene thus represents a novel growth factor which may be useful in tissue regeneration, and also represents a means for potentially inhibiting cell transformation via *wnt* signaling.

The *Frzb* proteins and the protein *Cerberus* are examples of secreted proteins that purportedly inhibit *wnt* signaling (Brown, *Curr. Opinion Cell Biol.*, 10:182-187 [1998]).

PCT WO 98/35043, published 13 August 1998 describes human SDF-5 proteins which are purportedly useful in regulating the binding of *wnt* polypeptides to their receptors.

PCT WO 98/23730, published 4 June 1998, describes transfecting tumors cells with *wnt-5a* to purportedly decrease tumorigenicity. *Wnt-5a* purportedly is an antagonist of other *wnts*.

In view of the devastating effects of cancer, there is a need in the art to identify additional genes that may serve as antagonists of proteins involved in cell transformation.

Accordingly, it is an object of this invention to provide nucleic acid molecules and polypeptides that may be useful as anti-cancer compounds.

It is a further object to provide methods of altering the level of expression and/or activity of such polypeptides in the human body.

- 3 -

Other related objects will readily be apparent from a reading of this disclosure.

## SUMMARY OF THE INVENTION

5

In one embodiment, the present invention provides an isolated nucleic acid molecule encoding a biologically active DKR polypeptide selected from the group consisting of:

- 10 (a) the nucleic acid molecule comprising SEQ  
ID NO:1;
- (b) the nucleic acid molecule comprising SEQ  
ID NO:2;
- 15 (c) the nucleic acid molecule comprising SEQ  
ID NO:3;
- (d) the nucleic acid molecule comprising SEQ  
ID NO:4;
- 20 (e) the nucleic acid molecule comprising SEQ  
ID NO:5;
- (f) the nucleic acid molecule comprising SEQ  
ID NO:6;
- (g) the nucleic acid molecule comprising SEQ  
ID NO:7;
- 25 (h) the nucleic acid molecule comprising SEQ  
ID NO:75;
- (i) the nucleic acid molecule comprising SEQ  
30 ID NO:76;
- (j) the nucleic acid molecule comprising SEQ  
ID NO:77;
- 35 (k) the nucleic acid molecule comprising SEQ  
ID NO:78;

- 4 -

- (l) the nucleic acid molecule encoding the polypeptide of SEQ ID NO:8;
- (m) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:9;
- (n) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:10, or a biologically active fragment thereof;
- (o) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:11, or a biologically active fragment thereof;
- (p) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:12, or a biologically active fragment thereof;
- (q) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:13, or a biologically active fragment thereof;
- (r) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:14, or a biologically active fragment thereof;
- (s) a nucleic acid molecule that encodes a polypeptide that is at least 85 percent identical to the polypeptide of SEQ ID NOS: 10, 11, 12, 13, or 14;
- (t) a nucleic acid molecule that encodes a biologically active DKR polypeptide that has 1-100 amino acid substitutions and/or deletions as compared with the polypeptide of any of SEQ ID NOS:8, 9, 10, 11, 12, 13, or 14; and
- (u) a nucleic acid molecule that hybridizes under conditions of high stringency to any of (c), (d), (e), (f), (g), (h), (i), (k), (l), (m), (n), (o), (p), (q), (r), (s), and (t) above.

In another embodiment, the invention provides an isolated nucleic acid molecule that is the complement of any of the nucleic acid molecules above.

- 5 -

In yet another embodiment, the invention provides an isolated nucleic acid molecule encoding a biologically active DKR polypeptide selected from the group of: amino acids 16-350, 21-350, 22-350, 23-350, 33-350, or 42-350, 21-145, 40-145, 40-150, 45-145, 45-145, 145-290, 150-290, 300-350, or 310-350 of SEQ ID NO:9; amino acids 15-266, 24-266, or 32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or 34-359 of SEQ ID NO:12; and amino acids 19-224, 20-224, 21-224, or 22-224 of SEQ ID NO:14.

In other embodiments, the invention provides vectors comprising the nucleic acid molecules, and host cells comprising the vectors.

In still another embodiment, the invention provides a process for producing a biologically active DKR polypeptide comprising the steps of:

- (a) expressing a polypeptide encoded by any of the nucleic acid molecules herein in a suitable host; and
- (b) isolating the polypeptide.

In still one other embodiment, the invention provides a biologically active DKR polypeptide selected from the group consisting of:

- (a) the polypeptide of SEQ ID NO:8;
- (b) the polypeptide of SEQ ID NO:9;
- (c) the polypeptide of SEQ ID NO:10;
- (d) the polypeptide of SEQ ID NO:11;
- (e) the polypeptide of SEQ ID NO:12;
- (f) the polypeptide of SEQ ID NO:13;
- (g) the polypeptide of SEQ ID NO:14;
- (h) a polypeptide that has 1-100 amino acid substitutions or deletions as compared with the polypeptide of any of (a)-(g) above; and
- (i) a polypeptide that is at least 85 percent identical to any of the polypeptides of (c)-(h) above.

- 6 -

In still one other embodiment, the invention provides the following polypeptides: a polypeptide that is amino acids 16-350, 21-350, 22-350, 23-350, 33-350, or 42-350, 21-145, 40-145, 40-150, 45-145, 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, or 310-350 of Figure 9, a polypeptide that is amino acids 15, 266, 24-266, or 32-266 of Figure 10, a polypeptide that is amino acids 17-259, 26-259, or 34-259 of Figure 12, and a polypeptide that is amino acids 19-224, 20-224, 21-224, or 22-224 of Figure 14.

## BRIEF DESCRIPTION OF THE DRAWINGS

15 Figure 1 (SEQ ID NO:1) depicts the cDNA sequence of mouse DKR-3.

20 Figure 2 (SEQ ID NO:2) depicts the cDNA sequence of human DKR-3.

Figure 3 (SEQ ID NO:3) depicts the DNA sequence of human DKR-1.

25 Figure 4 (SEQ ID NO:4) depicts the cDNA sequence of mouse DKR-2.

Figure 5 (SEQ ID NO:5) depicts the cDNA sequence of human DKR-2.

30 Figure 6 (SEQ ID NO:6) depicts the cDNA sequence of human DKR-2a, a splice variant of the DKR-2 gene.

35 Figure 7 (SEQ ID NO:7) depicts the cDNA sequence of human DKR-4.



- 7 -

Figure 8 (SEQ ID NO:8) depicts the amino acid sequence of mouse DKR-3 as translated from the corresponding cDNA.

5           Figure 9 (SEQ ID NO:9) depicts the amino acid sequence of human DKR-3 as translated from the corresponding cDNA.

10           Figure 10 (SEQ ID NO:10) depicts the amino acid sequence of human DKR-1 as translated from the corresponding cDNA.

15           Figure 11 (SEQ ID NO:11) depicts the amino acid sequence of mouse DKR-2 as translated from the corresponding cDNA.

20           Figure 12 (SEQ ID NO:12) depicts the amino acid sequence of human DKR-2 as translated from the corresponding cDNA.

            Figure 13 (SEQ ID NO:13) depicts the amino acid sequence of human DKR-2a as translated from the corresponding cDNA.

25           Figure 14 (SEQ ID NO:14) depicts the amino acid sequence of human DKR-4 as translated from the corresponding cDNA.

30           Figures 15A-15D are photographs of Northern blots which were probed with human DKR-3. Figure 15A shows the transcript level of DKR-3 in various human normal (Lanes 1-2) and immortal (Lanes 3-4) cell lines, and in human estrogen receptor plus ("ER+"; Lanes 5-9) and estrogen receptor minus ("ER-"; Lanes 10-16) breast cancer cell lines. Figure 15B shows the transcript level of human DKR-3 in human normal lung cells (Lane

35

- 8 -

1), and in various human non-small cell lung cancer ("NSCLC"; Lanes 2-9) and small cell lung cancer ("SCLC"; Lanes 10-15) cell lines. Figure 15C shows the amount of transcript of human DKR-3 in five glioblastoma cell lines; three of these lines (SNB-19, U-87MG, and U-373MG) are capable of forming tumors in nude mice, while the other two lines (Hs 683 and A 172) are not. Figure 15D shows the transcript level of human DKR-3 in human immortal (non-cancerous) and normal cervical cells, and in human cervical cancer cell lines (indicated as "tumor cells").

Figure 16 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

Figure 17 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

Figure 18 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

Figure 19 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

Figure 20 is a photograph of SDS gel electrophoresis. The contents of the lanes are set forth in the Examples herein.

Figure 21 is a photograph of a Western blot. Contents of the Lanes are indicated in the Examples herein.

- 9 -

Figure 22 (SEQ ID NO:75) is a nucleic acid sequence of human DKR-1 with codons optimized for expression in *E. coli*.

5           Figure 23 (SEQ ID NO:76) is a nucleic acid sequence of human DKR-2 with codons optimized for expression in *E. coli*.

10           Figure 24 (SEQ ID NO:77) is a nucleic acid sequence of human DKR-3 with codons optimized for expression in *E. coli*.

15           Figure 25 (SEQ ID NO:78) is a nucleic acid sequence of human DKR-4 with codons optimized for expression in *E. coli*.

#### DETAILED DESCRIPTION OF THE INVENTION

20           Included in the scope of this invention are *DKR* polypeptides such as the polypeptides of SEQ ID Nos:8-14, and related biologically active polypeptide fragments, variants, and derivatives thereof.

25           Also included within the scope of the present invention are nucleic acid molecules that encode *DKR* polypeptides such as the nucleic acid molecules of SEQ ID Nos:1-7.

30           Additionally included within the scope of the present invention are non-human mammals such as mice, rats, rabbits, goats, or sheep in which the gene (or genes) encoding a native *DKR* polypeptide has (have) been disrupted ("knocked out") such that the level of expression of this gene or genes is (are) significantly decreased or completely abolished. Such mammals may be prepared using techniques and methods such as those described in U.S. Patent No. 5,557,032. The present

35

- 10 -

invention further includes non-human mammals such as mice, rats, rabbits, goats, or sheep in which the gene (or genes) encoding *DKR* polypeptides in which either the native form of the gene(s) for that mammal or a heterologous *DKR* polypeptide gene(s) is (are) over expressed by the mammal, thereby creating a "transgenic" mammal. Such transgenic mammals may be prepared using well known methods such as those described in U.S. Patent No 5,489,743 and PCT patent application no. WO94/28122, published 8 December 1994. The present invention further includes non-human mammals in which the promoter for one or more of the *DKR* polypeptides of the present invention is either activated or inactivated (using homologous recombination methods as described below) to alter the level of expression of one or more of the native *DKR* polypeptides.

The *DKR* polypeptides of the present invention are expected to have utility as anti-cancer therapeutics for those cancers such as mammary tumors, stem cell tumors, or other cancers in which the *wnt* and/or sonic hedgehog (*shh*) signal transduction pathways are activated. Specific *wnt* members can transform mammary tissue (Hunter, supra) and are abnormally expressed in many human tumors (Huguet, *Cancer Res.*, 54:2615-2621 [1994]; Dale, *Cancer Res.*, 56:4320-4323 [1996]; see also PCT WO 97/39357). Such activity is expected in view of data presented herein in which the level of *DKR-3* transcript is decreased or not detectable at all in many cancer cell lines as compared to similar normal cell lines. Further, such activity is expected in view of the relationship of the genes and polypeptides of the present invention to the gene *dickkopf-1* (which, as mentioned above, is purportedly a potent antagonist of *wnt-8*). *DKR-1*, a novel gene of the present invention, is a human

- 11 -

ortholog of *dkk-1*. DKR-2, DKR-3, and DKR-4, all novel genes of the present invention, are each related to DKR-1 by their cysteine pattern. In particular, these DKR polypeptides may be of use for treatment of stem cell tumors, mammary tumors, and other cancers in which *wnt* genes are expressed, and in cancers where *wnt* and/or *shh* signaling is activated..

The DKR polypeptides of the present invention may also be administered as agents that can induce and/or enhance tissue differentiation, such as bone formation, cartilage formation, muscle tissue formation, nerve tissue formation, and hematopoietic cell formation. Such activities are expected in view of the fact that a) *Xenopus dkk-1* purportedly promotes head induction, heart formation, and differentiation of the developing CNS (Glinka, supra); and b) certain *wnt* polypeptides appear to function in embryo development (Cadigan, *Genes and Devel.*, 11:3286-3305 [1997]), specifically development of the pituitary (Treier, *Genes and Devel.*, 12:1691-1704 [1998]), myogenesis (Munsterberg et al., *Genes and Devel.*, 9:2911-2922 [1995]), osteogenesis (PCT WO 95/17416; PCT WO98/16641), kidney development (Stark et al., *Nature* 372:679-683 [1994]), development of the CNS (Dickinson et al., *Development*, 120:1453-1471 [1994]), and hematopoiesis (PCT WO 98/06747). Thus, addition of certain DKR polypeptides in such cell cultures or tissues may serve to modify the activity of various *wnt* polypeptides in cellular differentiation processes.

The DKR polypeptides herein may be used in either an *in vivo* manner or an *ex vivo* manner for such applications. For example, one or more of the DKR polypeptides of the present invention may be added to a culture of cartilage tissue or nerve tissue, or hematopoietic stem cells, either alone, or in combination with other growth factors and/or other

- 12 -

tissue differentiation factors, so as to induce or enhance the regeneration of such tissues.

Alternatively, such *DKR* polypeptides of the present invention may, for example, be injected directly into a joint in need of cartilage, into the spinal cord where the cord has been damaged, into damaged brain tissue, or into bone marrow to enhance hematopoiesis.

The term "*DKR* polypeptides" as used herein refers to any protein or polypeptide having the properties described herein for *DKR* polypeptides. The *DKR* polypeptides may or may not have amino terminal methionines, depending on the manner in which they are prepared. By way of illustration, *DKR* polypeptides refers to (1) a biologically active polypeptide encoded by any of the *DKR* polypeptides nucleic acid molecules as defined in any of items (a)-(f) below; (2) naturally occurring allelic variants and synthetic variants of any of *DKR* polypeptide in which one or more amino acid substitutions, deletions, and/or insertions are present as compared to the *DKR* polypeptides of SEQ ID NOs:8-14, and/or (3) biologically active polypeptides, or fragments or variants thereof, that have been chemically modified.

As used herein, the term "*DKR* polypeptide fragment" refers to a peptide or polypeptide that is less than the full length amino acid sequence of a naturally occurring *DKR* polypeptide but has the biological activity of any of the *DKR* polypeptides provided herein. Such a fragment may be truncated at the amino terminus, the carboxy terminus, and/or internally (such as by natural splicing), and may be a variant or a derivative of any of the *DKR* polypeptides. Such *DKR* polypeptides fragments may be prepared with or without an amino terminal methionine. In addition, *DKR* polypeptides fragments can be naturally occurring fragments such as *DKR* polypeptide splice variants (SEQ

- 13 -

ID NO:13), other splice variants, and fragments resulting from naturally occurring *in vivo* protease activity. Preferred DKR polypeptide fragments include amino acids 16-350, 21-350, 22-350, 23-350, 33-350, 42-350, 21-145, 40-145, 40-150, 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, and 310-350, of SEQ ID NO:9; amino acids 15-266, 24-266, or 32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or 34-359 of SEQ ID NO:12; and amino acids 19-224, 20-224, 21-224, or 22-224 of SEQ ID NO:14.

As used herein, the term "*DKR polypeptide variants*" refers to *DKR* polypeptides whose amino acid sequences contain one or more amino acid sequence substitutions, deletions, and/or insertions as compared to the *DKR* polypeptides amino acid sequences set forth in SEQ ID NOS:8-14. Such *DKR* polypeptides variants can be prepared from the corresponding *DKR* polypeptides nucleic acid molecule variants, which have a DNA sequence that varies accordingly from the DNA sequences for wild type *DKR* polypeptides as set forth in SEQ ID NOS:7-14. Preferred variants of the human *DKR* polypeptides include alanine substitutions at one or more of amino acid positions. Other preferred substitutions include conservative substitutions at the amino acid positions indicated in the Examples herein, as well as those encoded by *DKR* nucleic acid molecules as described below.

As used herein, the term "*DKR polypeptide derivatives*" refers to *DKR* polypeptides, variants, or fragments thereof, that have been chemically modified, as for example, by addition of one or more polyethylene glycol molecules, sugars, phosphates, and/or other such molecules, where the molecule or molecules are not naturally attached to wild-type *DKR* polypeptides.

As used herein, the terms "biologically active *DKR* polypeptides", "biologically active *DKR*

- 14 -

polypeptide fragments", "biologically active DKR polypeptide variants", and "biologically active DKR polypeptide derivatives" refer to DKR polypeptides that have the ability to decrease cancer cell proliferation in the Anchorage Independent Growth Assay of Example 12 herein, or in the *In Vivo* Tumor Assay of Example 13 herein, or in both assays.

As used herein, the term "DKR polypeptide nucleic acids" when used to describe a nucleic acid molecule refers to a nucleic acid molecule or fragment thereof that (a) has the nucleotide sequence as set forth in any of SEQ ID NOS:1-7; (b) has a nucleic acid sequence encoding a polypeptide that is at least 85 percent identical, but may be greater than 85 percent, i.e., 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent identical to the polypeptide encoded by any of SEQ ID NOS:10-14; (c) is a naturally occurring allelic variant or alternate splice variant of (a) or (b); (d) is a nucleic acid variant of (a)-(c) produced as provided for herein; (e) has a sequence that is complementary to (a)-(d); (f) hybridizes to any of (a)-(e) under conditions of high stringency and/or (g) has a nucleic acid sequence encoding 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or up to 100 amino acid substitutions and/or deletions of any mature DKR polypeptide (i.e., a DKR polypeptide with its endogenous signal peptide removed).

Percent sequence identity can be determined by standard methods that are commonly used to compare the similarity in position of the amino acids of two polypeptides. By way of example, using a computer algorithm such as GAP (Genetic Computer Group, University of Wisconsin, Madison, WI), the two polypeptides for which the percent sequence identity is



- 15 -

to be determined are aligned for optimal matching of their respective amino acids (the "matched span", as determined by the algorithm). A gap opening penalty (which is calculated as 3 X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect amino acid by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix (see Dayhoff et al., in: *Atlas of Protein Sequence and Structure*, vol. 5, supp.3 [1978] for the PAM250 comparison matrix; see Henikoff et al., *Proc. Natl. Acad. Sci USA*, 89:10915-10919 [1992] for the BLOSUM 62 comparison matrix) is also used by the algorithm. The percent identity is then calculated by the algorithm by determining the percent identity as follows:

- 16 -

Total number of identical matches  
in the matched span

$$\frac{[\text{length of the longer sequence} \\ \text{within the matched span}] + \\ [\text{number of gaps introduced into} \\ \text{the longer sequence in order to} \\ \text{align the two sequences}]}{\quad} \times 100$$

Polypeptides that are at least 85 percent identical will typically have one or more amino acid substitutions, deletions, and/or insertions as compared with any of the wild type *DKR* polypeptides. Usually, the substitutions of the native residue will be either alanine, or a conservative amino acid so as to have little or no effect on the overall net charge, polarity, or hydrophobicity of the protein. Conservative substitutions are set forth in Table I below.

- 17 -

Table I  
Conservative Amino Acid Substitutions

Basic:	arginine
	lysine
	histidine
Acidic:	glutamic acid
	aspartic acid
Uncharged Polar:	glutamine
	asparagine
	serine
	threonine
	tyrosine
Non-Polar:	phenylalanine
	tryptophan
	cysteine
	glycine
	alanine
	valine
	proline
	methionine
	leucine
	isoleucine

5           The term " conditions of high stringency"  
refers to hybridization and washing under conditions  
that permit binding of a nucleic acid molecule used for  
screening, such as an oligonucleotide probe or cDNA  
molecule probe, to highly homologous sequences. An  
10   exemplary high stringency wash solution is 0.2 X SSC  
and 0.1 percent SDS used at a temperature of between  
50°C-65°C.

          Where oligonucleotide probes are used to  
screen cDNA or genomic libraries, one of the following  
15   two high stringency solution may be used. The first of  
these is 6 X SSC with 0.05 percent sodium pyrophosphate

- 18 -

at a temperature of 35°C-62°C, depending on the length of the oligonucleotide probe. For example, 14 base pair probes are washed at 35-40°C, 17 base pair probes are washed at 45-50°C, 20 base pair probes are washed at 52-57°C, and 23 base pair probes are washed at 57-63°C. The temperature can be increased 2-3°C where the background non-specific binding appears high. A second high stringency solution utilizes tetramethylammonium chloride (TMAC) for washing oligonucleotide probes. One stringent washing solution is 3 M TMAC, 50 mM Tris-HCl, pH 8.0, and 0.2 percent SDS. The washing temperature using this solution is a function of the length of the probe. For example, a 17 base pair probe is washed at about 45-50°C.

As used herein, the terms "effective amount" and "therapeutically effective amount" refer to the amount of a *DKR* polypeptide necessary to support one or more biological activities of the *DKR* polypeptides as set forth above.

A full-length *DKR* polypeptide or fragment thereof can be prepared using well known recombinant DNA technology methods such as those set forth in Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY [1989]) and/or Ausubel et al., eds., (*Current Protocols in Molecular Biology*, Green Publishers Inc. and Wiley and Sons, NY [1994]). A gene or cDNA encoding a *DKR* polypeptide or fragment thereof may be obtained for example by screening a genomic or cDNA library, or by PCR amplification. Probes or primers useful for screening the library can be generated based on sequence information for other known genes or gene fragments from the same or a related family of genes, such as, for example, conserved motifs found in other *DKR* polypeptides such as the cysteine pattern. In addition, where a gene encoding *DKR*

- 19 -

polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify homologous genes from other species. The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the *DKR* gene. Typically, conditions of high stringency will be employed for screening to minimize the number of false positives obtained from the screen.

Another means to prepare a gene encoding a *DKR* polypeptide or fragment thereof is to employ chemical synthesis using methods well known to the skilled artisan such as those described by Engels *et al.* (*Angew. Chem. Intl. Ed.*, 28:716-734 [1989]). These methods include, *inter alia*, the phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, the DNA encoding the *DKR* polypeptide will be several hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form the full length *DKR* polypeptide. Usually, the DNA fragment encoding the amino terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of the *DKR* polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell.

In some cases, it may be desirable to prepare nucleic acid and/or amino acid variants of the naturally occurring *DKR* polypeptides. Nucleic acid variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate methods, where the primer(s) have the desired point mutations (see Sambrook *et al.*, *supra*, and Ausubel *et*

- 20 -

al., *supra*, for descriptions of mutagenesis techniques). Chemical synthesis using methods described by Engels et al., *supra*, may also be used to prepare such variants. Other methods known to the skilled artisan may be used as well. Preferred nucleic acid variants are those containing nucleotide substitutions accounting for codon preference in the host cell that is to be used to produce the *DKR* polypeptide(s). Such "codon optimization" can be determined via computer algorithms which incorporate codon frequency tables such as "Ecohigh. Cod" for codon preference of highly expressed bacterial genes as provided by the University of Wisconsin Package Version 9.0, Genetics Computer Group, Madison, WI. Other useful codon frequency tables include "Celegans\_high.cod", "Celegans\_low.cod", "Drosophila\_high.cod", "Human\_high.cod", "Maize\_high.cod", and "Yeast\_high.cod". Other preferred variants are those encoding conservative amino acid changes as described above (e.g., wherein the charge or polarity of the naturally occurring amino acid side chain is not altered substantially by substitution with a different amino acid) as compared to wild type, and/or those designed to either generate a novel glycosylation and/or phosphorylation site(s), or those designed to delete an existing glycosylation and/or phosphorylation site(s).

The gene, cDNA, or fragment thereof encoding the *DKR* polypeptide can be inserted into an appropriate expression or amplification vector using standard ligation techniques. The vector is typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). The gene, cDNA or fragment thereof encoding the *DKR* polypeptide may be

- 21 -

amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or eukaryotic host cells. Selection of the host cell will depend in part on whether the *DKR* polypeptide or fragment thereof is to be glycosylated and/or phosphorylated. If so, yeast, insect, or mammalian host cells are preferable.

Typically, the vectors used in any of the host cells will contain 5' flanking sequence (also referred to as a "promoter") and other regulatory elements as well such as an enhancer(s), an origin of replication element, a transcriptional termination element, a complete intron sequence containing a donor and acceptor splice site, a signal peptide sequence, a ribosome binding site element, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Each of these elements is discussed below. Optionally, the vector may contain a "tag" sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of the *DKR* polypeptide coding sequence; the oligonucleotide molecule encodes polyHis (such as hexaHis), or other "tag" such as FLAG, HA (hemagglutinin Influenza virus) or *myc* for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as means for affinity purification of the *DKR* polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified *DKR* polypeptide by various means such as using certain peptidases.

The human immunoglobulin hinge and Fc region could be fused at either the N-terminus or C-terminus of the *DKR* polypeptides by one skilled in the art. The

- 22 -

subsequent Fc-fusion protein could be purified by use of a Protein A affinity column. Fc is known to exhibit a long pharmacokinetic half-life *in vivo* and proteins fused to Fc have been found to exhibit a substantially greater half-life *in vivo* than the unfused counterpart. Also, fusion to the Fc region allows for dimerization/multimerization of the molecule that may be useful for the bioactivity of some molecules.

The 5' flanking sequence may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of 5' flanking sequences from more than one source), synthetic, or it may be the native *DKR* polypeptides gene 5' flanking sequence. As such, the source of the 5' flanking sequence may be any unicellular prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the 5' flanking sequence is functional in, and can be activated by, the host cell machinery.

The 5' flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, 5' flanking sequences useful herein other than the *DKR* gene flanking sequence will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of the 5' flanking sequence may be known. Here, the 5' flanking sequence may be synthesized using the methods described above for nucleic acid synthesis or cloning.

Where all or only a portion of the 5' flanking sequence is known, it may be obtained using PCR and/or by screening a genomic library with suitable



- 23 -

oligonucleotide and/or 5' flanking sequence fragments from the same or another species.

Where the 5' flanking sequence is not known, a fragment of DNA containing a 5' flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion using one or more carefully selected enzymes to isolate the proper DNA fragment. After digestion, the desired fragment may be isolated by agarose gel purification, Qiagen® column or other methods known to the skilled artisan. Selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

The origin of replication element is typically a part of prokaryotic expression vectors purchased commercially, and aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for optimal expression of the *DKR* polypeptide. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector.

The transcription termination element is typically located 3' of the end of the *DKR* polypeptide coding sequence and serves to terminate transcription of the *DKR* polypeptide. Usually, the transcription termination element in prokaryotic cells is a G-C rich fragment followed by a poly T sequence. While the element is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described above.

A selectable marker gene element encodes a protein necessary for the survival and growth of a host

- 24 -

cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic host cells, (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene.

The ribosome binding element, commonly called the Shine-Dalgarno sequence (prokaryotes) or the Kozak sequence (eukaryotes), is usually necessary for translation initiation of mRNA. The element is typically located 3' to the promoter and 5' to the coding sequence of the *DKR* polypeptide to be synthesized. The Shine-Dalgarno sequence is varied but is typically a polypurine (i.e., having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth above and used in a prokaryotic vector.

In those cases where it is desirable for *DKR* polypeptide to be secreted from the host cell, a signal sequence may be used to direct the *DKR* polypeptide out of the host cell where it is synthesized, and the carboxy-terminal part of the protein may be deleted in order to prevent membrane anchoring. Typically, the signal sequence is positioned in the coding region of the *DKR* gene or cDNA, or directly at the 5' end of the *DKR* gene coding region. Many signal sequences have been identified, and any of them that are functional in the selected host cell may be used in conjunction with the *DKR* gene or cDNA. Therefore, the signal sequence may be homologous or heterologous to the *DKR* gene or cDNA, and may be homologous or heterologous to the *DKR*

- 25 -

polypeptides gene or cDNA. Additionally, the signal sequence may be chemically synthesized using methods set forth above.

5 In most cases, secretion of the polypeptide from the host cell via the presence of a signal peptide will result in the removal of the amino terminal methionine from the polypeptide.

10 In many cases, transcription of the *DKR* gene or cDNA is increased by the presence of one or more introns in the vector; this is particularly true where the *DKR* polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within the *DKR* gene, especially where the gene used is a full length genomic  
15 sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron(s) may be obtained from another source. The position of the intron with respect to the 5' flanking sequence and the *DKR* gene is generally  
20 important, as the intron must be transcribed to be effective. As such, where the *DKR* gene inserted into the expression vector is a cDNA molecule, the preferred position for the intron is 3' to the transcription start site, and 5' to the polyA  
25 transcription termination sequence. Preferably for *DKR* cDNA, the intron or introns will be located on one side or the other (i.e., 5' or 3') of the cDNA such that it does not interrupt the this coding sequence. Any intron from any source, including any viral,  
30 prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this invention, provided that it is compatible with the host cell(s) into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the  
35 vector.

- 26 -

Where one or more of the elements set forth above are not already present in the vector to be used, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the  
5 elements are well known to the skilled artisan and are comparable to the methods set forth above (i.e., synthesis of the DNA, library screening, and the like).

The final vectors used to practice this invention are typically constructed from a starting  
10 vectors such as a commercially available vector. Such vectors may or may not contain some of the elements to be included in the completed vector. If none of the desired elements are present in the starting vector, each element may be individually ligated into the  
15 vector by cutting the vector with the appropriate restriction endonuclease(s) such that the ends of the element to be ligated in and the ends of the vector are compatible for ligation. In some cases, it may be necessary to "blunt" the ends to be ligated together in  
20 order to obtain a satisfactory ligation. Blunting is accomplished by first filling in "sticky ends" using Klenow DNA polymerase or T4 DNA polymerase in the presence of all four nucleotides. This procedure is well known in the art and is described for example in  
25 Sambrook et al., *supra*.

Alternatively, two or more of the elements to be inserted into the vector may first be ligated together (if they are to be positioned adjacent to each other) and then ligated into the vector.

30 One other method for constructing the vector to conduct all ligations of the various elements simultaneously in one reaction mixture. Here, many nonsense or nonfunctional vectors will be generated due to improper ligation or insertion of the elements,  
35 however the functional vector may be identified and selected by restriction endonuclease digestion.

- 27 -

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, *inter alia*, pCRII, PCR3, and pCDNA3.1 (Invitrogen Company, San Diego, CA), pBSII (Stratagene Company, La Jolla, CA), pET15b (Novagen, Madison, WI), pGEX (Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII; Invitrogen), and pFastBacDual (Gibco/BRL, Grand Island, NY).

After the vector has been constructed and a nucleic acid molecule encoding full length or truncated DKR polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression.

Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast cell, an insect cell, or a vertebrate cell). The host cell, when cultured under appropriate conditions, can synthesize DKR polypeptide which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). After collection, the DKR polypeptide can be purified using methods such as molecular sieve chromatography, affinity chromatography, and the like.

Selection of the host cell for DKR polypeptide production will depend in part on whether the DKR polypeptide is to be glycosylated or phosphorylated (in which case eukaryotic host cells are preferred), and the manner in which the host cell is able to "fold" the protein into its native tertiary structure (e.g., proper orientation of disulfide bridges, etc.) such that biologically active protein is prepared by the DKR polypeptide that has biological

- 28 -

activity, the *DKR* polypeptide may be "folded" after synthesis using appropriate chemical conditions as discussed below.

Suitable cells or cell lines may be mammalian cells, such as Chinese hamster ovary cells (CHO), human embryonic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. Other suitable mammalian cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines.

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 $\alpha$ , DH10, and MC1061) are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas* spp., other *Bacillus* spp., *Streptomyces* spp., and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for expression of the polypeptides of the present invention.

- 29 -

Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described for example in Kitts *et al.* (*Biotechniques*, 14:810-817 [1993]), Lucklow (*Curr. Opin. Biotechnol.*, 4:564-572 [1993]) and Lucklow *et al.* (*J. Virol.*, 67:4566-4579 [1993]). Preferred insect cells are Sf-9 and Hi5 (Invitrogen, Carlsbad, CA).

Insertion (also referred to as "transformation" or "transfection") of the vector into the selected host cell may be accomplished using such methods as calcium chloride, electroporation, microinjection, lipofection or the DEAE-dextran method. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook *et al.*, *supra*.

The host cells containing the vector (*i.e.*, transformed or transfected) may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing *E. coli* cells are for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells are RPMI 1640, MEM, DMEM, all of which may be supplemented with serum and/or growth factors as required by the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

Typically, an antibiotic or other compound useful for selective growth of the transformed cells only is added as a supplement to the media. The compound to be used will be dictated by the selectable

- 30 -

marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin.

5 The amount of *DKR* polypeptide produced in the host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, HPLC separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

If the *DKR* polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. 15 Polypeptides prepared in this way will typically not possess an amino terminal methionine, as it is removed during secretion from the cell. If however, the *DKR* polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus 20 (for eukaryotic host cells) or in the cytosol (for gram negative bacteria host cells) and may have an amino terminal methionine.

For *DKR* polypeptide situated in the host cell cytoplasm and/or nucleus, the host cells are typically 25 first disrupted mechanically or with detergent to release the intra-cellular contents into a buffered solution. *DKR* polypeptide can then be isolated from this solution.

Purification of *DKR* polypeptide from solution 30 can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (*DKR* polypeptide/hexaHis) or other small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or myc 35 (Invitrogen, Carlsbad, CA) at either its carboxyl or amino terminus, it may essentially be purified in a



- 31 -

one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag or for the polypeptide directly (i.e., a monoclonal antibody specifically recognizing DKR polypeptide). For example, polyhistidine binds with great affinity and specificity to nickel, thus an affinity column of nickel (such as the Qiagen® nickel columns) can be used for purification of DKR polypeptide/polyHis. (See for example, Ausubel et al., eds., *Current Protocols in Molecular Biology*, Section 10.11.8, John Wiley & Sons, New York [1993]).

Where the DKR polypeptide is prepared without a tag attached, and no antibodies are available, other well known procedures for purification can be used. Such procedures include, without limitation, ion exchange chromatography, molecular sieve chromatography, HPLC, native gel electrophoresis in combination with gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific). In some cases, two or more of these techniques may be combined to achieve increased purity.

If it is anticipated that the DKR polypeptide will be found primarily intracellularly, the intracellular material (including inclusion bodies for gram-negative bacteria) can be extracted from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

If the DKR polypeptide has formed inclusion bodies in the cytosol, the inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The

- 32 -

pellet material can then be treated at pH extremes or with chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The DKR polypeptide in its now soluble form can then be analyzed using gel electrophoresis, immunoprecipitation or the like. If it is desired to isolate the DKR polypeptide, isolation may be accomplished using standard methods such as those set forth below and in Marston et al. (*Meth. Enz.*, 182:264-275 [1990]). In some cases, the DKR polypeptide may not be biologically active upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure and generating disulfide linkages, can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization but usually at a lower concentration and is not necessarily the same chaotrope as used for the solubilization. In most cases the refolding/oxidation solution will also contain a reducing agent or the reducing agent plus its' oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein's cysteine bridge(s). Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, dithiothreitol (DTT)/dithiane DTT, 2-mercaptoethanol (BME)/dithio-b(ME). In many

instances a cosolvent is necessary to increase the efficiency of the refolding and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, and arginine.

If *DKR* polypeptide inclusion bodies are not formed to a significant degree in the host cell, the *DKR* polypeptide will be found primarily in the supernatant after centrifugation of the cell homogenate, and the *DKR* polypeptide can be isolated from the supernatant using methods such as those set forth below.

In those situations where it is preferable to partially or completely isolate the *DKR* polypeptide, purification can be accomplished using standard methods well known to the skilled artisan. Such methods include, without limitation, separation by electrophoresis followed by electroelution, various types of chromatography (immunoaffinity, molecular sieve, and/or ion exchange), and/or high pressure liquid chromatography. In some cases, it may be preferable to use more than one of these methods for complete purification.

In addition to preparing and purifying *DKR* polypeptide using recombinant DNA techniques, the *DKR* polypeptides, fragments, and/or derivatives thereof may be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art such as those set forth by Merrifield et al., (*J. Am. Chem. Soc.*, 85:2149 [1963]), Houghten et al. (*Proc Natl Acad. Sci. USA*, 82:5132 [1985]), and Stewart and Young (*Solid Phase Peptide Synthesis*, Pierce Chemical Co., Rockford, IL [1984]). Such polypeptides may be synthesized with or without a methionine on the amino terminus. Chemically synthesized *DKR*

- 34 -

polypeptides or fragments may be oxidized using methods set forth in these references to form disulfide bridges. The *DKR* polypeptides or fragments are expected to have biological activity comparable to *DKR* polypeptides produced recombinantly or purified from natural sources, and thus may be used interchangeably with recombinant or natural *DKR* polypeptide.

Chemically modified *DKR* polypeptide compositions in which *DKR* polypeptide is linked to a polymer are included within the scope of the present invention. The polymer selected is typically water soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. The polymer selected is usually modified to have a single reactive group, such as an active ester for acylation or an aldehyde for alkylation, so that the degree of polymerization may be controlled as provided for in the present methods. The polymer may be of any molecular weight, and may be branched or unbranched. Included within the scope of *DKR* polypeptide polymers is a mixture of polymers. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

The water soluble polymer or mixture thereof may be selected from the group consisting of, for example, polyethylene glycol (PEG), monomethoxy-polyethylene glycol, dextran, cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (e.g., glycerol) and polyvinyl alcohol.

For the acylation reactions, the polymer(s) selected should have a single reactive ester group.

- 35 -

For reductive alkylation, the polymer(s) selected should have a single reactive aldehyde group. A preferred reactive aldehyde is polyethylene glycol propionaldehyde, which is water stable, or mono C1-C10 alkoxy or aryloxy derivatives thereof (see U.S. Patent No. 5,252,714).

Pegylation of *DKR* polypeptides may be carried out by any of the pegylation reactions known in the art, as described for example in the following references: *Focus on Growth Factors* 3: 4-10 (1992); EP 0 154 316; and EP 0 401 384. Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described below.

A particularly preferred water-soluble polymer for use herein is polyethylene glycol, abbreviated PEG. As used herein, polyethylene glycol is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono-(C1-C10) alkoxy- or aryloxy-polyethylene glycol.

In general, chemical derivatization may be performed under any suitable conditions used to react a biologically active substance with an activated polymer molecule. Methods for preparing pegylated *DKR* polypeptides will generally comprise the steps of (a) reacting the polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby *DKR* polypeptide becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined based on known parameters and the desired result. For example, the larger the ratio of PEG: protein, the greater the percentage of poly-pegylated product.

- 36 -

Generally, conditions which may be alleviated or modulated by administration of the present polymer/polypeptides include those described herein for *DKR* polypeptides molecules. However, the polymer/ *DKR* polypeptides molecules disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased half-life, as compared to the non-derivatized molecules.

The *DKR* polypeptides, fragments thereof, variants, and derivatives, may be employed alone, together, or in combination with other pharmaceutical compositions. The *DKR* polypeptides, fragments, variants, and derivatives may be used in combination with cytokines, growth factors, antibiotics, anti-inflammatory, and/or chemotherapeutic agents as is appropriate for the indication being treated.

*DKR* nucleic acid molecules, fragments, and/or derivatives that do not themselves encode polypeptides that are active in activity assays may be useful as hybridization probes in diagnostic assays to test, either qualitatively or quantitatively, for the presence of *DKR* DNA or corresponding RNA in mammalian tissue or bodily fluid samples.

*DKR* polypeptide fragments, variants, and/or derivatives that are not themselves active in activity assays may be useful for preparing antibodies that recognize *DKR* polypeptides.

The *DKR* polypeptides, fragments, variants, and/or derivatives may be used to prepare antibodies using standard methods. Thus, antibodies that react with the *DKR* polypeptides, as well as reactive fragments of such antibodies, are also contemplated as within the scope of the present invention. The antibodies may be polyclonal, monoclonal, recombinant, chimeric, single-chain and/or bispecific. Typically,

- 37 -

the antibody or fragment thereof will either be of human origin, or will be "humanized", i.e., prepared so as to prevent or minimize an immune reaction to the antibody when administered to a patient. The antibody  
5 fragment may be any fragment that is reactive with *DKR* polypeptides of the present invention, such as, Fab, Fab', etc. Also provided by this invention are the hybridomas generated by presenting any *DKR* polypeptide or fragments thereof as an antigen to a selected  
10 mammal, followed by fusing cells (e.g., spleen cells) of the mammal with certain cancer cells to create immortalized cell lines by known techniques. The methods employed to generate such cell lines and antibodies directed against all or portions of a human  
15 *DKR* polypeptide of the present invention are also encompassed by this invention.

The antibodies may be used therapeutically, such as to inhibit binding of the *DKR* polypeptide to its binding partner. The antibodies may further be  
20 used for *in vivo* and *in vitro* diagnostic purposes, such as in labeled form to detect the presence of *DKR* polypeptide in a body fluid or cell sample.

Preferred antibodies are human antibodies, either polyclonal or monoclonal.  
25

#### Therapeutic Compositions and Administration

Therapeutic compositions of *DKR* polypeptides are within the scope of the present invention. Such compositions may comprise a therapeutically effective  
30 amount of the polypeptide or fragments, variants, or derivatives in admixture with a pharmaceutically acceptable carrier. The carrier material may be water for injection, preferably supplemented with other materials common in solutions for administration to  
35 mammals. Typically, a *DKR* polypeptide therapeutic

- 38 -

compound will be administered in the form of a composition comprising purified polypeptide, fragment, variant, or derivative in conjunction with one or more physiologically acceptable carriers, excipients, or diluents. Neutral buffered saline or saline mixed with serum albumin are exemplary appropriate carriers. Preferably, the product is formulated as a lyophilizate using appropriate excipients (e.g., sucrose). Other standard carriers, diluents, and excipients may be included as desired. Other exemplary compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor.

The DKR polypeptide compositions can be administered parenterally. Alternatively, the compositions may be administered intravenously or subcutaneously. When systemically administered, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such pharmaceutically acceptable protein solutions, with due regard to pH, isotonicity, stability and the like, is within the skill of the art.

Therapeutic formulations of DKR polypeptide compositions useful for practicing the present invention may be prepared for storage by mixing the selected composition having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (*Remington's Pharmaceutical Sciences*, 18th Edition, A.R. Gennaro, ed., Mack Publishing Company [1990]) in the form of a lyophilized cake or an aqueous solution. Acceptable carriers, excipients or stabilizers are nontoxic to recipients and are preferably inert at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or other organic acids;



- 39 -

antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

An effective amount of the *DKR* polypeptide composition(s) to be employed therapeutically will depend, for example, upon the therapeutic objectives such as the indication for which the *DKR* polypeptide is being used, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage may range from about 0.1  $\mu\text{g/kg}$  to up to 100 mg/kg or more, depending on the factors mentioned above. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of *DKR* polypeptide) over time, or as a continuous infusion via implantation device or catheter.

As further studies are conducted, information will emerge regarding appropriate dosage levels for treatment of various conditions in various patients, and the ordinary skilled worker, considering the therapeutic context, the type of disorder under

- 40 -

treatment, the age and general health of the recipient, will be able to ascertain proper dosing.

The *DKR* polypeptide composition to be used for *in vivo* administration must be sterile. This is  
5 readily accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using these methods may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral  
10 administration ordinarily will be stored in lyophilized form or in solution.

Therapeutic compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a  
15 stopper pierceable by a hypodermic injection needle.

The route of administration of the composition is in accord with known methods, e.g. oral, injection or infusion by intravenous, intraperitoneal, intracerebral (intraparenchymal),  
20 intracerebroventricular, intramuscular, intraocular, intraarterial, or intralesional routes, or by sustained release systems or implantation device which may optionally involve the use of a catheter. Where desired, the compositions may be administered  
25 continuously by infusion, bolus injection or by implantation device.

Alternatively or additionally, the composition may be administered locally via implantation into the affected area of a membrane,  
30 sponge, or other appropriate material on to which *DKR* polypeptide has been absorbed.

Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of *DKR* polypeptide may be directly  
35 through the device via bolus, or via continuous

- 41 -

administration, or via catheter using continuous infusion.

DKR polypeptide may be administered in a sustained release formulation or preparation. Suitable examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides (U.S. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al., *Biopolymers*, 22: 547-556 [1983]), poly (2-hydroxyethyl-methacrylate) (Langer et al., *J. Biomed. Mater. Res.*, 15: 167-277 [1981] and Langer, *Chem. Tech.*, 12: 98-105 [1982]), ethylene vinyl acetate (Langer et al., *supra*) or poly-D(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also may include liposomes, which can be prepared by any of several methods known in the art (e.g., Eppstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688-3692 [1985]; EP 36,676; EP 88,046; EP 143,949).

In some cases, it may be desirable to use DKR polypeptide compositions in an *ex vivo* manner. Here, cells, tissues, or organs that have been removed from the patient are exposed to DKR polypeptide compositions after which the cells, tissues and/or organs are subsequently implanted back into the patient.

In other cases, DKR polypeptide may be delivered through implanting into patients certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the polypeptides, fragments, variants, or derivatives. Such cells may be animal or human cells, and may be derived from the patient's own tissue or from another source, either human or non-human. Optionally, the cells may be immortalized. However, in order to decrease the chance of an immunological

- 42 -

response, it is preferred that the cells be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or  
5 membranes that allow release of the protein product(s) but prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

Methods used for membrane encapsulation of  
10 cells are familiar to the skilled artisan, and preparation of encapsulated cells and their implantation in patients may be accomplished without undue experimentation. See, e.g., U.S Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for  
15 encapsulating living cells is described in PCT WO 91/10425 (Aebischer et al.). Techniques for formulating a variety of other sustained or controlled delivery means, such as liposome carriers, bio-erodible particles or beads, are also known to those in the art,  
20 and are described, for example, in U.S. Patent No. 5,653,975 (Baetge et al., CytoTherapeutics, Inc.). The cells, with or without encapsulation, may be implanted into suitable body tissues or organs of the patient.

As discussed above, it may be desirable to  
25 treat isolated cell populations such as stem cells, lymphocytes, red blood cells, chondrocytes, neurons, and the like with one or more *DKR* polypeptides, variants, derivatives and/or fragments. This can be accomplished by exposing the isolated cells to the  
30 polypeptide, variant, derivative, or fragment directly, where it is in a form that is permeable to the cell membrane. Alternatively, gene therapy can be employed as described below.

One manner in which gene therapy can be  
35 applied is to use the *DKR* gene (either genomic DNA, cDNA, and/or synthetic DNA encoding a *DKR* polypeptide,

- 43 -

or a fragment, variant, or derivative thereof) which may be operably linked to a constitutive or inducible promoter to form a "gene therapy DNA construct". The promoter may be homologous or heterologous to the endogenous *DKR* gene, provided that it is active in the cell or tissue type into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include, as required, DNA molecules designed for site-specific integration (e.g., endogenous flanking sequences useful for homologous recombination), tissue-specific promoter, enhancer(s) or silencer(s), DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (as, for example, for cell targeting) cell-specific internalization factors, and transcription factors to enhance expression by a vector as well as factors to enable vector manufacture.

This gene therapy DNA construct can then be introduced into the patient's cells (either *ex vivo* or *in vivo*). One means for introducing the gene therapy DNA construct is via viral vectors. Suitable viral vectors typically used in gene therapy for delivery of gene therapy DNA constructs include, without limitation, adenovirus, adeno-associated virus, herpes simplex virus, lentivirus, papilloma virus, and retrovirus vectors. Some of these vectors, such as retroviral vectors, will deliver the gene therapy DNA construct to the chromosomal DNA of the patient's cells, and the gene therapy DNA construct can integrate into the chromosomal DNA; other vectors will function as episomes and the gene therapy DNA construct will remain in the cytoplasm. The use of gene therapy vectors is described, for example, in U.S. Patent Nos. 5,672,344 (30 September 1997; Kelly et al., University

- 44 -

of Michigan), 5,399,346 (21 March 1995; Anderson et al., U.S. Dept. Health and Human Services), 5,631,236 (20 May 1997; Woo et al., Baylor College of Medicine), and 5,635,399 (3 June 1997; Kriegler et al., Chiron Corp.).

Alternative means to deliver gene therapy DNA constructs to a patient's cells without the use of viral vectors include, without limitation, liposome-mediated transfer, direct injection of naked DNA, receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation, and microparticle bombardment (e.g., "gene gun"). See U.S. Patent Nos. 4,970,154 (13 November 1990; Chang, Baylor College of Medicine), WO 96/40958 (19 December 1996; Smith et al., Baylor College of Medicine) 5,679,559 (21 October 1997; Kim et al., University of Utah) 5,676,954 (14 October 1997; Brigham, Vanderbilt University), and 5,593,875 (14 January 1997; Wurm et al., Genentech).

Another means to increase endogenous *DKR* polypeptide expression in a cell via gene therapy is to insert one or more enhancer elements into the *DKR* polypeptide promoter, where the enhancer element(s) can serve to increase transcriptional activity of the *DKR* polypeptides gene. The enhancer element(s) used will be selected based on the tissue in which one desires to activate the gene(s); enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a *DKR* polypeptide is to be "turned on" in T-cells, the *lck* promoter enhancer element may be used. Here, the functional portion of the transcriptional element to be added may be inserted into a fragment of DNA containing the *DKR* polypeptide promoter (and optionally, vector, 5' and/or 3' flanking sequence, etc.) using standard cloning techniques. This construct, known as a "homologous recombination

- 45 -

construct" can then be introduced into the desired cells either *ex vivo* or *in vivo*.

Gene therapy can be used to decrease *DKR* polypeptide expression by modifying the nucleotide  
5 sequence of the endogenous promoter(s). Such modification is typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the *DKR* gene(s) selected for inactivation can be engineered to  
10 remove and/or replace pieces of the promoter that regulate transcription. Here, the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter  
15 activity thereby repressing transcription of the corresponding *DKR* gene. Deletion of the TATA box or transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the *DKR*  
20 polypeptide promoter(s) (from the same or a related species as the *DKR* gene(s) to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are mutated via substitution, deletion and/or insertion of one or more  
25 nucleotides such that the TATA box and/or activator binding site has decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' flanking regions of the promoter segment that has been  
30 modified, may be introduced into the appropriate cells (either *ex vivo* or *in vivo*) either directly or via a viral vector as described above. Typically, integration of the construct into the genomic DNA of  
35 the cells will be via homologous recombination, where the 5' and 3' flanking DNA sequences in the promoter

- 46 -

construct can serve to help integrate the modified promoter region via hybridization to the endogenous chromosomal DNA.

Other gene therapy methods may also be employed where it is desirable to inhibit one or more *DKR* polypeptides. For example, antisense DNA or RNA molecules, which have a sequence that is complementary to at least a portion of the selected *DKR* polypeptide gene(s) can be introduced into the cell. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected *DKR* gene. When the antisense molecule then hybridizes to the corresponding *DKR* polypeptides mRNA, translation of this mRNA is prevented.

Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more of the *DKR* polypeptides. In this situation, the DNA encoding a mutant full length or truncated polypeptide of each selected *DKR* polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described above. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

Samples of the *E coli* cell lines GM121 and GM94 have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA, USA on September 22, 1998 as accession numbers 202173 and 202174, respectively.

The following examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.



- 47 -

## EXAMPLES

Example 1: Cloning of the Mouse DKR-3 Gene

5 About 120 adult mice with an average body weight of about 18 grams were each injected intraperitoneally with a kainate solution (prepared as a stock solution of about 1 mg/ml kainate in sterile PBS) at a dose of about 25 mg kainate per kilogram body weight. About six hours after injection, the mice were sacrificed, and the hippocampus was dissected from each mouse. Total RNA was extracted from hippocampal tissue using the Trizol method (Gibco BRL, Grand Island, NY). The poly(A+) mRNA fraction was isolated from total RNA using Message Maker (Gibco BRL, Grand Island, NY) according to the manufacturer's recommended procedure. Hippocampal tissue was also obtained from control mice (which received an injection of PBS only), and poly(a+) mRNA was obtained from this tissue as well using the same procedures.

Two random primed cDNA libraries were prepared; one from the kainate-treated and one from the control poly (A+) mRNA using the Superscript® plasmid system (Gibco BRL, Gaithersburg, MD). A random cDNA primer containing an internal *NotI* restriction site was used to initiate first strand synthesis and had the following sequence:

30 GGAAGGAAAAAAGCGGCCGCAACANNNNNNNNNN (SEQ ID NO:15)

where N is A, G, C, or T.

Both first strand cDNA synthesis and second strand cDNA synthesis were performed according to the manufacturer's recommended protocol. After second

35

- 48 -

strand synthesis, the reaction products were extracted with phenol:chloroform:isoamyl alcohol (in a volume ratio of 25:24:1), followed by ethanol precipitation. The double strand cDNA products were ligated using  
5 standard ligation procedures to the following double stranded oligonucleotide adapter (obtained from Gibco BRL, Grand Island, NY):

TCGACCCACGCGTCCG (SEQ ID NO:16)  
10. GGGTGCGCAGGC (SEQ ID NO:17)

After ligation, the cDNA was digested to completion with *NotI*, and size fractionated on a 1  
15 percent agarose gel. The cDNA products between about 250 and 800 base pairs were selected and purified from the gel using the Qiagen® gel extraction kit (Qiagen, Chatsworth, CA). The purified cDNA products were directionally ligated into the vector pYY41L (American  
20 Type Culture Collection, "ATCC"; 10801 University Blvd., Manassas, VA, USA; accession number 209636) which had been previously digested with *NotI* and *SalI*. The ligated cDNA was then introduced into electrocompetent ElectroMax® DH108 *E. coli* cells  
25 (Gibco-BRL, Grand Island, NY) via standard electroporation techniques. The library was then titered by a serial dilution of the transformation cell mixture.

About one million primary clones were divided  
30 into 20 pools (50,000 clones each pool) and each pool was plated on 245mm x 245 mm square plate containing MR2001 medium (MacConnel Research, San Diego, CA) and about 60 ug/ml carbonocillin. After incubation overnight at 37C, the colonies were scraped off the  
35 plate in about 20 ml SOC (SOC contains about 2 percent

- 49 -

Bactotryptone, 0.5 percent yeast extract, 10 mM sodium chloride, 2.5 mM potassium chloride, and 10 mM magnesium sulfate) and were pelleted by centrifugation at about 6000 rpm for about 10 minutes. The plasmids  
5 were then recovered from the cells using Qiagen® maxi prep columns (Qiagen, Chatsworth, CA) according to the protocol suggested by the manufacturer.

About two hundred and fifty thousand clones (50 ug total plasmids/10 ug from each pool) were used  
10 to transform yeast strain YPH499 (ATCC accession number 90834) and an amylase-based signal trap assay was conducted as follows (see co-pending U.S.S.N. 09/026,959 filed 20 February 1998 for a detailed description of this technique). Around 1000  
15 transformants were plated on a single starch-containing selection plate (15 cm diameter with a medium containing about 0.6 percent yeast nitrogen base, 2 percent glucose, 0.1 percent CAA, 1.0 X trp dropout solution, 0.7 percent potato starch azure, and 1.5  
20 percent agarose). The plates were incubated at about 30C for 4-5 days until full development of halos was observed. The colonies in the center of the halo were picked and restreaked on a fresh plate to form single colonies. The single colonies with halos were then  
25 picked and arrayed into 96 well microtiter plates containing about 100 ul of water per well, thereby generating the "yeast colony solutions".

About ten microliters of each well of each yeast colony solution was used as template to recover  
30 the cDNA fragment from that colony through PCR. Therefore, ninety-six PCR reactions were independently performed using PCR-Ready Beads® (96 well format, Amersham-Pharmacia Biotech, Piscataway, NJ) and the following oligonucleotides according to the  
35 manufacturer's protocol:

- 50 -

ACTAGCTCCAGTGATCTC (SEQ ID NO:18)

CGTCATTGTTCTCGTTCC (SEQ ID NO:19)

5

PCR was conducted using a Perkin-Elmer 9600 thermocycler with the following cycle conditions: 94C for 10 minutes followed by 35 cycles of 94C for 30 seconds, 55C for 30 seconds and 72C for 1 minute, after which a final extension cycle of 72C for 10 minutes was conducted. Most PCR reactions contained a single PCR product. The amplified cDNA products were purified using the Qiagen® PCR purification kit (Qiagen, Chatsworth, CA). These products were sequenced on an Applied Biosystems 373A automated DNA sequencer using the following oligonucleotide primer:

GCTATACCAAGCATACAATC (SEQ ID NO:35)

20

Taq dye-terminator sequencing reactions (Applied Biosystems, Foster City, CA) were conducted following the manufacturer's recommended procedures.

Each PCR fragment was translated in all six possible ways to identify those fragments which (1) had a potential signal peptide in the same direction as reporter gene; (2) had a stop codon(s) upstream of the putative methionine translation start site; and (3) appeared to lack a transmembrane domain.

One clone that met these criteria, termed "ymrs2-00009-c4", was selected for further analysis. This clone contained 5' sequence, including a putative signal sequence, but was lacking 3' sequence.

To obtain the 3' sequence of this clone, a 3' RACE reaction was performed using as a template pool

35

- 51 -

number 4 from the YmHK2 cDNA library. This YmHK2 library was prepared as follows: First strand cDNA synthesis was performed using about 2 micrograms of the RNA obtained from the hippocampus of the kainate  
5 treated mice and about 1 ug of *Not I* primer-adapter having the following sequence:

GACTAGTTCTAGATCGCGAGCGGCCGCCCTTTTTTTTTTTTTTTT (SEQ ID  
NO:42)

10

Both the first strand and second strand cDNA synthesis reactions were performed using the Superscript® plasmid system (Gibco BRL, Grand Island, NY). After second strand synthesis, the double  
15 stranded cDNA products were ligated into the double stranded adapters of SEQ ID NOs:16 and 17.

After ligation, the cDNA was digested to completion with *Not I*, and size fractionated on a 0.8 percent agarose gel. The cDNA products larger than  
20 about 800 base pairs were selected and purified from the gel using the Qiagen® gel extraction kit (Qiagen, Chatsworth, CA). The purified cDNA products were directly ligated into *Sal I* and *Not I* digested pSport® vector (Gibco BRL, Grand Island, NY).

25 The ligated cDNA products were then introduced into electrocompetent *E. coli* cells called ElectroMax® DH10B (Gibco BRL, Grand Island, NY). The library was then titered.

About twelve million primary clones were  
30 obtained, and expanded into about 250 ml of LB containing about 100 ug/ml ampicillin. After overnight incubation at 37C, the plasmids were recovered using the Qiagen® maxi-prep kit (Qiagen, Chatsworth, CA).

- 52 -

About 20 ng of the plasmid library were used to transform the ElectroMax® DH10B electrocompetent *E. coli* cells using standard electroporation techniques. About two million transformants were divided into 40  
5 pools (containing approximately 50,000 plasmids/pool). Each pool was then expanded into about 3 ml of LB medium containing about 100 ug/ml ampicillin. After overnight incubation at 37C, the plasmids were recovered using the Qiagen® mini-prep kit. The DNA  
10 from each pool were then stored at about minus 20C for future use.

The 3' RACE reaction was performed using about 1.5 ng of pool #4 of the YmHK2 library as a template, and using the Advantage® cDNA PCR kit  
15 (Clontech, Palo Alto, CA) with the following oligonucleotides:

CCAGCTGCTCTGTGGCAGCCCAG (SEQ ID NO:20)  
20 CCCAGTCACGACGTTGTAAAACGACGGCC (SEQ ID NO:21)

The reaction was conducted in a standard thermocycler (Perkin-Elmer 9600) for thirty five cycles under the following conditions: 94 C for 1 minute; 94 C  
25 for 5 seconds, and 72 C for 5 minutes. This was followed by a final extension at 72C for 10 minutes. About one microliter of the reaction products was diluted to 50 ul using TE buffer (10 mM TRIS pH 8.0 and 1 mM EDTA).

30 To enrich the RACE reaction for the gene of interest, a nested PCR reaction was conducted using about five microliters of the TE solution (containing the RACE reaction products as described in the preceding paragraph) together with the following  
35 oligonucleotides:

- 53 -

AACATGCAGCGGCTCGGGGG

(SEQ ID NO:22)

GGTGACACTATAGAAGAGCTATGACGTCGC

(SEQ ID NO:23)

5

The nested PCR reaction was incubated in a thermocycler (Perkin-Elmer 9600) using the following protocol: 94C for one minute; five cycles of 94C for 5 seconds followed by 72C for 5 minutes; five cycles of 94C for five seconds, followed by 70C for 5 minutes; and 20-25 cycles of 94C for 5 seconds followed by 68C for 5 minutes. After this PCR, the 3' RACE products and the nested PCR products were analyzed using standard agarose gel electrophoresis.

15

A PCR product of about 3.3 kb was identified from the nested PCR reaction. This fragment was purified using Qiagen® Gel Extraction Kit (Qiagen, Chatsworth, CA) and ligated into the vector pCRII-TOPO (Invitrogen, Carlsbad, CA) according to the procedures recommended by the manufacturer. After ligation, the products were transformed into One Shot® *E. coli* cells (Invitrogen, Carlsbad, CA) and plated on a LB (Luria broth) plate containing about 100 ug/ml ampicillin and about 1.6 mg X-gal. After overnight incubation at 37C, 12 white colonies and one blue colony were selected, and screened using PCR-Ready Beads® (Amersham-Pharmacia Biotech, Piscataway, NJ) according to the manufacturer's recommended protocol using oligonucleotide SEQ ID NO:20 together with the following primer:

30

GTGCTGAGTGTCTTCCATCAGC

(SEQ ID NO:24)

Two colonies were picked that had yielded PCR products of the expected size of about 192 base pairs.

35

- 54 -

These colonies were inoculated into about 3 ml of LB medium containing about 100 ug/ml ampicillin, and were incubated at 37C. The cultures were placed on a shaker for about 16 hours, and the plasmids were recovered using Qiagen® mini prep columns (Qiagen, Chatsworth, CA) according to the manufacturer's protocol. Plasmid DNA was then sequenced as described above.

A contiguous stretch of DNA of about 3366 nucleotides was assembled by combining the sequence of clone ymrs2-00009-c4 (containing 5' sequence) together with the nested PCR fragment of 3.3 kb containing 3' sequence. Within this contiguous sequence is an open reading frame of 349 amino acids. The nucleotide sequence of this novel mouse gene, referred to as DKR-3, is set forth in Figure 1. The putative amino acid sequence, as translated from the DNA sequence, is set forth in Figure 8

A BLAST search of the Genbank database using the amino acid sequence of DKR-3 revealed that this open reading frame has homology to a gene known as human rig-like 7-1 mRNA (Genbank accession number AF034208; see also Ligon *et al.*, *J NeuroVirology*, 4:217-226 [1998]). DKR-3 also has homology to the gene for chicken lens fiber protein clfest4 (Genbank accession number D26311); the overall identity to this protein is about 50 percent with the highest homology in the middle of the protein.

#### Example 2: Cloning of the Human DKR-3 Gene

Mouse DKR-3 DNA can be used to search a public EST database for human homologs, resulting in the identification of the following Genbank accession numbers:



- 55 -

AA628979  
AA349552  
AA633061  
AA351624  
5 W61032  
T30923  
AA683017  
AA324686  
T08793  
10 T31076  
R14945  
AA226979  
W45085  
AA424460  
15 R58671  
R57834  
AF034208

These EST sequences were analyzed and  
20 assembled to create a putative sequence for human DKR-  
3. Based on this putative sequence, two  
oligonulceotides were designed for use in PCR in an  
attempt to clone the human DKR-3 gene. The sequence of  
these oligonucleotides is:

25 GAGATGCAGCGGCTTGGGGCCACCC (SEQ ID NO:25)

GCCTGGTCAGCCCACGCCTAAAG (SEQ ID NO:26)

30 PCR was performed using the Advantage® cDNA  
PCR kit (Clontech, Palo Alto, CA) together with human  
fetal brain Quick-Clone® cDNA (Clontech). PCR was  
conducted in a thermocycler (Perkin-Elmer 9600) under  
the following cycle conditions: 94C for 2 minute; 94C  
35 for 30 seconds, and 72C for 2 minutes. Thirty-five

- 56 -

cycles were conducted after which samples were treated at 72C for 10 minutes. A single fragment of about 1150 base pairs was visible when the PCR products were visualized on a 1 percent agarose gel. This fragment

5 was purified using the Qiagen® Gel Extraction Kit (Qiagen, Chatsworth, CA) and ligated into the vector pCRII-TOPO (Invitrogen, Carlsbad, CA). After ligation, the products were transformed into One Shoot *E. coli*® (Invitrogen, Carlsbad, CA) and plated on a LB plate

10 containing about 100 ug/ml ampicillin and about 1.6 mg X-gal. After overnight incubation at 37C, 2 white colonies were picked and inoculated into about 3 ml of LB medium containing about 100 ug/ml ampicillin. The cultures were kept on a shaker at about 37C for about

15 16 hours. The plasmids were isolated using Qiagen® mini-prep columns (Qiagen, Chatsworth, CA) according to the manufacturer's recommended protocol, and the inserts were then sequenced using methods described above.

20 The cloned fragment is 1141 bp in length and contains an open reading frame of 350 amino acids. The nucleotide sequence is set forth in Figure 2, and the putative amino acid sequence, as translated from the DNA sequence, is set forth in Figure 9. This amino

25 acid sequence is about 80 percent identical to the mouse DKR-3 gene. In addition, human DKR-3 is identical to the human rig-like protein fragment described by Lignon et al., *supra* between amino acids 157 and 308 of DKR-3. Significantly, the rig-like

30 protein has an amino terminal start corresponding to amino acid 156 of DKR-3. Rig-like does not appear to be a secreted protein, and the carboxy terminal region of rig-like protein has no homology to human DKR-3. Just as for mouse DKR-3, human DKR-3 is about 54

35 percent identical to the chicken lens fiber protein

- 57 -

cldest. Human DKR-3 appears to be secreted, with a signal peptide cleavage site after either amino acid 20 or 21. Other potential cleavage sites (due to signal peptides or to other endogenous processing sites are after amino acid 16, 22, 32, and/or 41). There appear to be N-linked glycosylation sites at amino acids 96, 106, 121, and 204, which would render them preferable sites for generating substitution mutants. Human DKR-3 and mouse DKR-3 amino acid sequences differ at amino acid positions 6, 7, 11, 24, 27, 29, 30, 32, 33, 39, 81, 89, 93, 99, 101, 103, 109, 113, 115, 123, 126, 142, 156, 157, 162, 165, 173, 175, 191, 197, 198, 201, 203, 245, 247, 259, 283, 287, 292, 294, 295, 296, 298, 299, 304, 310, 311, 312, 314, 315, 329, 330, 334, 335, 336, 339, 340, 341, 342, 343, 345, and 347 (all with respect to the human DKR-3 sequence), which renders these positions preferable for generating human DKR-3 substitution or deletion variants. Based on computer analysis of the amino acid sequence of DKR-3, significant regions of the molecule include the span from about amino acids 21-145 (a potential alpha-helical region and region of potential N-linked glycosylation) such as for example amino acids 21-145, 40-145, 40-150, 45-145, and 45-150, the span from about amino acids 145-350, such as, for example 145-290, 145-300, and 145-350, and the span from about amino acids 300-350 (a second potential alpha-helical region), such as for example amino acids 310-350. Such regions would be suitable fragments of full length DKR-3.

Northern blot analysis was conducted to assess the tissue specific expression of human DKR-3. A probe for use in Northern blot analysis was prepared by PCR of human fetal brain Quick-Clone® cDNA (Clontech, Palo Alto, CA) using the following oligonucleotides:

- 58 -

CCTGCTGCTGGCGGCGGCGGTCCCCACGGC

(SEQ ID NO:27)

GCCTGGTCAGCCCACGCCTAAAG

(SEQ ID NO:28)

5

The PCR reaction was conducted in a thermocycler (Perkin-Elmer 9600). PCR conditions were: 94C for 2 minute; 94C for 30 seconds, and 72C for 2 and 1/2 minutes. Thirty-five cycles were conducted followed by a final extension treatment at 72C for 10 minutes. PCR products were run on a one percent agarose gel, and a band of about 1100 bp was gel purified using the Qiagen gel extraction kit (Qiagen®, Chatsworth, CA), cloned into the vector CRII-TOPO (Invitrogen, Carlsbad, CA) and sequenced to confirm that the band contained the human DKR-3 open reading frame minus the amino terminal 10 amino acids.

About twenty-five nanograms of this probe was denatured by heating to about 100C for about 5 minutes, followed by placing on ice, and then radioactively labeled with alpha-32P-dCTP using the Rediprime® labeling kit (Amersham, Arlington Heights, IL) and following the manufacturer's instructions. A human multiple tissue Northern blot was purchased (Clontech, Palo Alto, CA) and was first prehybridized in about 5 ml of Clontech Express® hybridization buffer at about 68C for 30-60 minutes. After prehybridization, the labeled probe was added to the solution and allowed to hybridize for about 60 minutes. After hybridization, the blot was first washed with 2xSSC plus 0.05 percent SDS at room temperature for about 30 minutes, then washed with 0.1xSSC plus 0.1 percent SDS at about 65C for about 30 minutes. The blot was dried briefly and then exposed to a Phosphorimager screen (Molecular Dynamics, Sunnyvale, CA). After overnight exposure,

- 59 -

the image of the blot was analyzed on a Storm 820 machine (Molecular Dynamics, Sunnyvale, CA) with Imagequat software (Molecular Dynamics, Sunnyvale, CA).

5 The size of the human DKR-3 RNA transcript is about 2.6 kb. The results of the Northern blot analysis indicate that human DKR-3 is highly expressed in adult heart and brain, although weak expression in placenta, adult lung, skeletal muscle, kidney, and pancreas is also apparent. A second smaller transcript is apparent  
10 in adult pancreas, and could result from degradation of the full length transcript.

To evaluate the role of this gene in cancer, a variety of human cancer cell lines were analyzed for the presence or absence of DKR-3 RNA transcript.

15 The glioblastoma cell lines Hs 683; A 172; SNB-19; U-87MG; and U-373MG are all from ATCC, and cultured in the media recommended by ATCC.

Normal human mammary epithelial cells (NMECs) derived from reduction mammoplasties were purchased  
20 from Clonetics Corp. (San Diego, CA) and the Corriel Institute (Camden, N.J.). The immortalized breast epithelial cell line MCF-10 and the ER+ cell line MCF-7 can be obtained from the American Type Culture Collection. The ER+ BT20T cells were provided by Dr.  
25 K. Keyomarsi (N.Y. State Dept. of Health). Immortalized 184A1 and other breast cancer cells including T47-D, ZR75-1, and BT474, MDA-MB-157, MDA-MB-231, MDA-MB-361, MDA-MB-453, MD-MBA-468, HS578T and SKBr3 were all obtained from the American Type Culture  
30 Collection (10801 University Blvd., Manassas, VA).

NMECs, 184A1 and MCF10 cells were cultured in a modified DME/F12 medium (Gibco/BRL, Grand Island, NY) supplemented with 10 mM Hepes, 2mM glutamine, 0.1 mM nonessential amino acids, 0.5 mM ethanolamine, 5 mg/ml  
35 transferrin, 1 mg/ml Bovine serum albumin, 5.0 ng/ml sodium selenite, 20 ng/ml triiodothyronine, 10 ng/ml

- 60 -

EGF, 5 µg/ml insulin and 0.5 µg/ml hydrocortisone (DMEM/F12C) (Ethier *et al*, *Cancer Letters*, 74:189-195 [1993]). The ER+ and ER+ breast cancer cells were cultured in Alpha or Richter improved minimal essential medium (MEM) (Gibco/BRL) supplemented with 10 mM Hepes, 2mM glutamine, 0.1 mM nonessential amino acids, 10 percent fetal bovine serum and 1 µg/ml insulin.

Normal human bronchial and cervical epithelial cells were purchased from Clonetics Corp. (San Diego, CA). Normal cervical epithelial cells were culture in KBM2 (Clonetics Corp. San Diego, CA) supplemented with 13 mg/ml bovine pituitary extract, 0.5 µg/ml hydrocortisone, 2 ng/ml EGF, 0.5 mg/ml epinephrine, 0.1 ng/ml retinoic acid, 5 µg/ml transferrin, 6.5 ng/ml triiodothyronine and 5 µg/ml insulin. Normal bronchial epithelial cells were cultured in BEBM (Clonetics Crop., San Diego, CA) supplemented with 0.5 mg/ ml hydrocortisone, 0.5 ng/ ml EGF, 0.5 µg/ml epinephrine, 10 µg/ml transferrin, 5 µg/ ml insulin, 0.1 ng/ml retinoic acid and 5.5 ng/ ml triiodthyronine.

The lung cancer cell lines H1299, H23, H358, H441, H460, H520, H522, H727, H146, H209, H446, H510A, H526, and H889 and the cervical cancer cells Caski, C-4-I, MS751, SiHa and C-33-A were all obtained from the American Type Culture Collection. The lung cancer cells were cultured in RPMI (MEM) (Gibco/BRL) supplemented with 10 mM Hepes, 2 mM glutamine, 0.1 mM nonessential amino acids and 10 percent fetal bovine serum (FBS). The cervical cancer cells were cultured in Earles MEM supplemented with 0.1 mM nonessential amino acids, 1 mM sodium pyruvate and 10 percent FBS. All cells were routinely screened for mycoplasma

- 61 -

contamination and maintained at about 37°C in an atmosphere of about 6.5 percent CO<sub>2</sub>.

Total RNA was prepared by lysing cell monolayers in guanidinium isothiocyanate and centrifuging over a 5.7 M CsCl cushion as described previously (Gudas, *Proc. Natl. Acad. Sci USA*, 85:4705-4709 [1988]). RNA (about 20 ug) was electrophoresed on denaturing formaldehyde gels, transferred to MagnaNT membranes (Micron Separations Inc., Westboro, MA) and cross-linked with UV irradiation.

The blots were prehybridized, probed, and washed under the same conditions as those set forth above for the tissue blot. The blots were dried briefly and then exposed to a Phosphorimager screen (Molecular Dynamics, Sunnyvale, CA). After overnight exposure, the image of the blot was analyzed on a Storm 820 machine with Imagequat software (both from Molecular Dynamics).

The results are shown in Figures 15A-15D. As can be seen in Figure 15A, expression of DKR-3 is decreased in most of the breast cancer cell lines as compared to the normal cell lines. Figure 15B indicates that DKR-3 expression is decreased in the non-small cell lung cancer cell lines, and in most of the small cell lung cancer cell lines as well. Figure 15C indicates that expression of DKR-3 is decreased in three glioblastoma cell lines (SNB-19, U-87MG, and U-373MG) that are capable of forming tumors in nude mice (the other two cell lines, Hs 683 and A 172 do not form tumors in nude mice). Figure 15D indicates that expression of DKR-3 is reduced in cervical cancer cell lines as compared to normal and immortalized cells.

- 62 -

Example 3: Cloning of the Human DKR-1 Gene

Human and mouse DKR-3 cDNA and amino acid sequences were used to search Genbank using the BLAST program in an attempt to identify DKR-3 related genes. A number of EST (expressed sequence tags) were found and were analyzed to determine whether the sequences overlapped. Using the following human EST accessions, a novel gene, termed DKR-1, was predicted.

10  
15  
20  
25  
30  
35

AA336797  
R27865  
W39690  
AA043027  
HUM517H04B  
AA143670  
W51876  
N94525  
AA641247  
AA137219  
AA115249  
AA031969  
AA136192  
AA032060  
AA035583  
AA207078  
AA371363  
AA037322  
AA088618  
W46873  
AA115337  
AA693679  
W30750  
H83554



- 63 -

PCR was conducted in an attempt to clone the full length gene, and the following two oligonucleotides were used for PCR:

5 CCCGGACCCTGACTCTGCAGCCG (SEQ ID NO:29)

GAGGAAAAATAGGCAGTGCAGCACC (SEQ ID NO:30)

PCR was performed using the Advantage® cDNA PCR kit (Clontech, Palo Alto, CA) containing the oligonucleotides listed above and human placenta Quick-Clone® cDNA (Clontech, Palo Alto, CA). The reaction was conducted according to the manufacturer's recommendations. Thirty-five cycles of PCR were conducted in a thermocycler (Perkin-Elmer 9600) under the following conditions: 94C for 2 minutes; 94C for 30 seconds, and 72C for 1-1/2 minutes, followed by a final extension at 72C for 10 minutes.

After cycling, the PCR products were analyzed on a one percent agarose gel. A single band of about 1200 base pairs in length was detected after agarose gel electrophoresis. This fragment was purified using the Qiagen® gel extraction kit (Qiagen, Chatsworth, CA) and ligated into the vector pCRII-TOPO (Invitrogen, Carlsbad, CA) using standard ligation procedures. After ligation, the products were transformed into One Shoot® competent *E. coli* cells according to the procedures recommended by manufacturer (Invitrogen, Carlsbad, CA). The transformed *E. coli* cells were plated on a LB plate containing about 100 ug/ml ampicillin and about 1.6 mg X-gal.

After overnight incubation at about 37C, two white colonies were picked and inoculated into about 3 ml of TB containing 100 ug/ml ampicillin. The culture was incubated at about 37C for about 16 hours, plasmids

- 64 -

were then recovered using Qiagen® mini-prep columns (Qiagen, Chatsworth, CA) and sequenced. Both colonies contained the same insert.

The insert is 1193 base pairs, and is referred to as human DKR-1. The sequence of this gene is set forth in Figure 3. This gene contains an open reading frame of 266 amino acids. The amino acid sequence is set forth in Figure 10. A stop codon is present upstream of the first methionine, indicating the first methionine is likely to be the amino terminus of the protein. Human DKR-1 has a predicted signal peptide with a predicted signal peptide cleavage site between amino acids 19 and 20.

The gene has about 80 percent homology to the mouse gene *dkk-1* (Glinka et al., supra), however the mouse *dkk-1* gene is 272 amino acids in length while human DKR-1 is 266 amino acids in length. Human DKR-1 differs from mouse *dkk-1* at amino acid positions 3, 4, 5, 7, 8, 10, 12, 13, 14, 15, 16, 17, 18, 19, 22, 23, 24, 29, 53, 55, 62, 66, 69, 77, 93, 98, 101, 105, 106, 123, 139, 140, 143, 144, 153, 155, 157, 158, 163, 164, 165, 169, 175, 178, 197, 224, and 244. In addition, the alignment of human DKR-1 and mouse *dkk-1* shows one gap in human DKR-1 between amino acids 37 and 38, and two gaps between 103 and 104, 146 and 147, and 165 and 166. Glinka et al. state on page 362 of their article that "Coordinates of *Xenopus* *dkk* family members have been deposited in Genbank with the following accession numbers...hdkk-1 AA207078." However, forward three frame translations of AA207078 by the inventors herein showed no homology to the published mouse and *Xenopus* *dkk-1* sequences, or to the human DKR-1 sequence, except in the 3' end of this accession, which exhibits a 95 percent identity to human DKR-1 from amino acids 81-179, indicating that AA207078 does not encode full

- 65 -

length human *dkk-1*. Significantly, AA207078 is missing amino acids 1-90 and 180-350 of human DKR-1 which includes the signal peptide and the second cysteine right domain respectively.

5  
Example 4: Cloning of the Mouse DKR-2 Gene

Genbank accession number AA265561 (a mouse sequence) has homology to both human DKR-1 and human DKR-3 at the amino acid level based primarily on its cysteine pattern.

To extend this EST sequence in both the 5' and 3' directions, the following oligonucleotides were designed:

15 GCCACAGTCCCCACCAAGGATCATC

(SEQ ID NO:31)

GATGATCCTTGGTGGGGACTGTGGC

(SEQ ID NO:32)

20 CTGCAAACCAGTGCTCCATCAGGG

(SEQ ID NO:33)

CCCTGATGGAGCACTGGTTTGCAG

(SEQ ID NO:34)

25 Separately, 5' RACE and 3' RACE reactions were performed according to the manufacturer's protocol using mouse heart Marathon-Ready® cDNA and the Advantage® cDNA PCR kit (both from Clontech, Palo Alto, CA) and using oligonucleotide SEQ ID NOS: 31 and 34. The RACE reactions were incubated in a thermocycler (Perkin-Elmer 9600) using the following cycling conditions: 94C for one minute; five cycles of 94C for 5 seconds followed by 72C for 5 minutes; five cycles of 94C for five seconds, followed by 70C for 5 minutes; and 20-25 cycles of 94C for 5 seconds followed by 68C for 5 minutes.

30

35

- 66 -

To enrich each RACE reaction for the desired product, about one microliter of each of the RACE PCR products was added together, and the mixture was diluted to about 50 ul using TE buffer. About five  
5 microliters of this solution were used to conduct nested PCR reactions. The Advantage® cDNA PCR kit (Clontech, Palo Alto, CA) and oligonucleotide SEQ ID NOs: 32 and 33 were used for the 5' and 3' nesting reactions, respectively. The nested PCR reactions were  
10 incubated in a thermocycler (Perkin-Elmer 9600) using the following program for thirty five cycles: 94C for 1 minute; 94C for 5 seconds; and 72C for 2 minutes. A final extension was then conducted at 72C for 10 minutes. The PCR products were analyzed using a one  
15 percent agarose gel.

Several fragments ranging from about 500 bp to about 1500 base pairs were obtained from the 5' nested PCR reaction, and two fragments of about 1900 bp and 450 bp were obtained from the 3' nested PCR  
20 reaction. These PCR products were purified using the Qiagen® PCR purification kit (Qiagen, Chatsworth, CA) and were then ligated into the vector pCRII-TOPO (Invitrogen). The ligation products were transformed into OneShot® *E. coli* cells (Invitrogen, Carlsbad,  
25 CA), and the cells were then plated on to two X-gal containing plates (one for each reaction) as described above.

Eight white colonies from each plate were picked and PCR selected via RACE reactions using the  
30 Clontech primer AP2 and the oligonucleotide SEQ ID NO:32 (for the 5' RACE) or the oligonucleotide SEQ ID NO:33 (for the 3' RACE). Three colonies from each plate that contained the correct size fragments were cultured, and the plasmids were isolated and sequenced  
35 using procedures described above.

- 67 -

Three clones, 9813302, 9813304 and 9813305 contained sequence which extended the EST sequence in the 5' direction. One clone, 9813308, contained sequence which extended the EST sequence in the 3' direction. A continuous sequence of 2678 base pairs was thus assembled using the sequence of clones 9813308, 9813304, and the EST AA265561. This full length DNA has been termed DKR-2, and the sequence is set forth in Figure 4. The corresponding amino acid sequence is set forth in Figure 11.

Within the amino acid sequence is an open reading frame of 259 amino acids. This protein has approximately 38 percent identity with mouse *dkk-1* at the amino acid level. Mouse DKR-2 has a predicted signal peptide with a signal peptide cleavage site between amino acids 33 and 34.

#### Example 5: Cloning of the Human DKR-2 Gene

The Genbank EST database was searched using the BLAST program with both DNA and amino acid sequences from human DKR-1 and human DKR-3, and one human EST, W55979, was identified that showed homology to both human DKR-1 and human DKR-3 at the amino acid level based on its cysteine pattern. W55979 is about 88 percent identical to mouse DKR-2 at the DNA level, and about 93 percent identical to mouse DKR-2 at the amino acid level.

A BLAST search of Genbank W55979 indicated that W55979 has homology to BAC clone number B284B3 (Genbank accession number AC003099). BAC clone B284B3 is 95129 base pairs in length. Three portions of W55979 are homologous to three different regions of BAC clone B284B3, indicating that human DKR-2 has at least three exons. A 3' sequence of 556 bp in length was assembled based on the sequences of both BAC clone

- 68 -

B284B3 and W55979, and it was determined that this sequence is the 3' portion of the human ortholog of mouse DKR-2. Within this 3' sequence of human DKR-2 is an open reading frame of 174 amino acids, and a stop codon is present after amino acid 174. This 3' sequence of human DKR-2 is about 97 percent identical to mouse DKR-2.

To obtain the 5' end sequence of human DKR-2, a 5' RACE reaction was performed using Clontech human heart Marathon-Ready® cDNA and the Advantage® cDNA PCR kit, together with oligonucleotide SEQ ID NO:34. The RACE reaction was performed according to the manufacturer's protocol. The 5' RACE reaction products were then subjected to nesting PCR to enrich for the 5' sequence using the Advantage® cDNA PCR kit and oligonucleotide SEQ ID NO:32. The PCR conditions for both the 5' RACE reaction and the nested PCR reaction were the same as those described in Example 4.

The nested PCR products were purified using the Qiagen® (Qiagen, Chatsworth, CA) PCR purification kit, and were ligated into the vector Zero-Blunt® (Invitrogen, San Diego, CA) according to the procedures recommended by the manufacturer. The ligation products were transformed into OneShot® *E. coli* cells which were then plated on X-gal containing plates as described above.

After overnight culturing, three white colonies were picked and were inoculated into about 3 ml of TB containing about 100 ug/ml ampicillin. The cultures were allowed to grow for about 16 hours, after which the plasmids were isolated using Qiagen® mini-prep columns (Qiagen, Chatsworth, CA) according to the manufacturer's protocol. The sequence of each insert was then obtained.

- 69 -

One of the 5'-RACE clones, termed 9812826, extended the human DKR-2 sequence 5'-terminally. A contiguous sequence of 1531 bp in length was assembled using this clone 9812826 together with the human DKR-2 3'-sequence. Within this contiguous sequence is an open reading frame of 259 amino acids. The human DKR-2 gene has a predicted signal peptide of about 33 amino acids, with a predicted cut site between amino acids 33 and 34, and is about 95 percent identical to mouse DKR-2 at the amino acid level. The amino acid positions that differ between human and mouse DKR-2 include (with respect to the numbering of the human sequence) 7, 12, 28, 48, 50, 58, 71, 102, 119, 170, 173, and 191, rendering these positions preferable for generating amino acid substitution or deletion variants.

An alternative spliced isoform of human DKR-2 was discovered when PCR was conducted using human heart Marathon-Ready® cDNA (Clontech, Palo Alto, CA) and the Advantage® cDNA PCR kit (Clontech, Palo Alto, CA) together with the following oligonucleotides:

GGGTTGAGGGAACACAATCTGCAAG

(SEQ ID NO:36)

GTCTGCAATTGATGATGTTCTCAATGG

(SEQ ID NO:37)

PCR was conducted using parameters set forth in the manufacturer's protocol. PCR products were analyzed by agarose gel electrophoresis, and two PCR products were obtained. The bands corresponding to these products were gel purified as described above, amplified and purified as described above, and then sequenced. One product corresponded to full length DKR-2, however, the other band corresponded to an isoform of DKR-2. This isoform has an open reading frame of 207 amino acids, and appears to be missing an

- 70 -

exon. This isoform is referred to as human DKR-2a. The DNA sequence of human DKR-2a is set forth in Figure 6, and the amino acid sequence as translated from the DNA is set forth in Figure 13.

5

Example 6: Cloning of the Human DKR-4 Gene

A human EST that showed significant homology to human DKR-1 and human DKR-3 on protein level was identified in Genbank. This sequence, Genbank accession number AA565546, has a cysteine pattern that is similar to that of human DKR-1 and human DKR-3.

A BLAST search of Genbank showed no human ESTs overlapping with AA565546. Therefore, to extend the EST sequence in the 5' direction, a 5' RACE reaction was performed using human heart Marathon-Ready® cDNA (Clontech, Palo Alto, CA) together with the Advantage® cDNA PCR kit (Clontech, Palo Alto, CA) and the following oligonucleotide:

20

CCAGGGCCACAGTCGCAACGCTGG (SEQ ID NO:38)

The RACE reaction was performed according to the protocol provided with the Advantage® kit. After 5' RACE, the products were nested to enrich for the desired 5' sequence using the Advantage® cDNA PCR kit according to the manufacturer's recommendations, together with the following oligonucleotide:

30 CTCCCTCTTGTCCTTCCTGCCTTG (SEQ ID NO:39)

After the nested PCR reaction, the products were purified using the Qiagen® PCR purification kit (Qiagen, Chatsworth, CA), ligated into the vector



- 71 -

pCRII-TOPO (Invitrogen, Carlsbad, CA), and transformed into OneShot® *E. coli* cells as described above. After transformation, the cells were plated on a LB plate containing about 100 ug/ml ampicillin and about 1.6 mg X-gal.

After overnight incubation at 37C, four white colonies were picked from the plate and were inoculated in about 3 ml TB containing about 100 ug/ml ampicillin. The cultures were incubated at about 37C for about 16 hours. The plasmids were then recovered using Qiagen® mini-prep columns (Qiagen, Chatsworth, CA) and sequenced.

Two clones, termed 9813563 and 9853564, were found to contain the 5' sequence of human DKR-4.

To obtain the 3' sequence of human DKR-4, a 3' RACE reaction was performed using human uterus Marathon-Ready® cDNA (Clontech, Palo Alto, CA) together with the Advantage® cDNA PCR kit (Clontech) and the following oligonucleotide:

CAAGGCAGGAAGGGACAAGAGGGAG (SEQ ID NO:40)

The 3' RACE reaction was performed according to the manufacturer's recommendations. After the RACE reaction, the products were nested using the Advantage® cDNA PCR kit and the following oligonucleotide:

CCAGCGTTGCGACTGTGGCCCTGG (SEQ ID NO:41)

The parameters for PCR were 94C for 1 minute followed by thirty five cycles of 94C for 5 seconds and then 72C for 2 minutes, after which a final extension of 70C for 10 minutes was conducted. After the nesting

- 72 -

reaction, the products were analyzed on a 1 percent agarose gel. A single band of about 1200 bp in length was observed. This band was purified from the gel using methods described above, and was then cloned into the vector pCR2.1-TOPO (Invitrogen, Carlsbad, CA) and sequenced. Sequence of this band indicated that it contained the 3' sequence of human DKR-4., and this sequence was assembled together with the 5' sequence (from clones 9813563 and 9853564) to generated the full length sequence of human DKR-4. This sequence is set forth in Figures 7 (DNA sequence) and 14 (translated amino acid sequence). The polypeptide is 224 amino acids in length and is about 34 percent identical to human DKR-1 at the amino acid sequence level.

Example 7: Expression of Human DKR-1 in Bacteria

PCR amplification employing the primer pairs and template described below were used to generate a recombinant form of human DKR-1. One primer of each pair introduces a TAA stop codon and a unique *Bam*HI site following the carboxy terminus of the gene. The other primer of each pair introduces a unique *Nde*I site, a N-terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, restriction digested, and inserted into the unique *Nde*I and *Bam*HI sites of vector pAMG21 (ATCC accession no. 98113) and transformed into the prototrophic *E. coli* host GM121 (deposited with the American Type Culture Collection on September 22, 1998 as accession number 202174). Other commonly used *E. coli* expression vectors and host cells are also suitable for expression by one skilled in the art. After transformation,

- 73 -

positive clones were selected and examined for expression of the recombinant gene product.

The construct pAMG21-human DKR-1-24-266 was engineered to be 244 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-His-Pro-Leu-Leu-Gly (SEQ ID NO:43)

10 Thr-Cys-Gln-Arg-His (SEQ ID NO:44)

The template used for PCR was human DKR-1 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

15 GTTCTCCTCATATGCATCCATTATTAGGCGTAAGTGCCACCTTGAACCTCGGTTCT  
CAAT (SEQ ID NO:45)

20 TACGCACTGGATCCTTAGTGTCTCTGACAAGTGTGAAG (SEQ ID NO:46)

Transformed *E. coli* strain GM121 containing pAMG21-human DKR-1-24-266 were grown in 2X YT media containing 20 micrograms/ml kanamycin at 30C until the culture reached an optical density of about 600 nm of about 0.5. Induction of DKR-1 protein expression was achieved by addition of *Vibrio fischeri* synthetic autoinducer to 100 ng/ml final and incubation of the culture at either 30 °C or 37 °C for about 9 hours further with shaking. In addition, as a uninduced control, for each culture no autoinducer was added to an aliquot of the culture, but the culture was also incubated for about 9 hours further at about 30C with shaking along with the induced cultures. After about 9 hours, the optical density of cultures were measured at 600 nm, an aliquot of cultures were examined by oil emersion microscopy at 1600X magnification, and

- 74 -

aliquots of cultures were pelleted by centrifugation. Bacterial pellets of cultures were processed for SDS-polyacrylamide gel electrophoresis on a 14 percent gel to examine levels of protein produced in crude lysates and for N-terminal sequencing confirmation of the recombinant gene product. The gel was stained with Coomassie blue.

The results are shown in the photo of Figure 16. Lane 1 contains molecular weight markers; Lanes 2 and 5 contain crude lysates of uninduced control cells incubated at 30°C; Lanes 3 and 6 are crude lysates of induced cells cultured at 30°C; Lanes 4 and 7 are crude lysates of induced cells cultured at 37°C. The arrow on the left of Lane 1 indicates the expected location of human DKR-1-24-266. As can be seen, large amounts of recombinant protein were observed in crude lysates of induced cultures at both 30°C and 37°C (Lanes 3 and 6, and 4 and 7). Microscopic analysis of bacterial cells revealed most cells contained at least one inclusion body, suggesting that at least some of the protein may be produced in the insoluble fraction of *E. coli*.

#### Example 8: Expression of DKR-2 in Bacteria

PCR amplification employing the primer pairs and templates described below were used to generate various forms of DKR-2. One primer of each pair introduces a TAA stop codon and a unique *Bam*HI site following the carboxy terminus of the gene. The other primer of each pair introduces a unique *Nde*I site, a N-terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, restriction digested, and inserted into the unique *Nde*I and *Bam*HI sites of vector pAMG21 (ATCC accession no.

- 75 -

98113) and transformed into either prototrophic *E. coli* host GM121 or GM94 (GM 94 was deposited with the ATCC on September 22, 1998 as accession number 202173). Other commonly used *E. coli* expression vectors and host cells are also suitable for expression. After transformation, positive clones were selected and examined for expression of the recombinant gene product.

The construct pAMG21-human DKR-2-26-259 was engineered to be 235 amino acids in length and have the following N-terminal and the following C-terminal amino acids, respectively:

Met-Ser-Gln-Ile-Gly-Ser (SEQ ID NO:47)

Val-Cys-Gln-Lys-Ile (SEQ ID NO:48).

The template used for PCR was human DKR-2 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct.

GTTCTCCTCATATGTCTCAAATTGGTAGTTCTCGTGCCAACTCAACTCCATCAAG (SEQ ID NO:49)

TACGCACTGGATCCTTAAATTTTCTGACACACATGGAGT (SEQ ID NO:50)

The construct pAMG21 mouse DKR-2-26-259 was engineered to be 235 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Ser-Gln-Leu-Gly-Ser (SEQ ID NO:51)

Val-Cys-Gln-Lys-Ile (SEQ ID NO:52)

- 76 -

The template used for PCR was mouse DKR-2 cDNA, and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct.

5 GTTCTCCTCATATGTCTCAATTAGGTAGCTCTCGTGCTAAACTCAACTCCATCAA  
GTCC (SEQ ID NO:53)

TACGCACTGGATCCTTAGATCTTCTGGCATAACATGGAGT (SEQ ID NO:54)

10 Transformed *E. coli* GM121 or GM94 containing  
either pAMG21-human DKR-2-26-259 or pAMG21-mouse  
DKR-2-26-259 plasmid were grown in 2X YT media  
containing 20 µg/ml kanamycin at 30 °C until the culture  
reached an optical density at 600 nm of about 0.5.  
15 Induction of DKR-2 protein expression was achieved by  
addition of *Vibrio fischeri* synthetic autoinducer to  
100 ng/ml final and incubation of the culture at either  
30C or 37C for about 5 or 9 hours further with shaking.  
In addition, as a uninduced control, for each culture,  
20 no autoinducer was added to an aliquot of the culture,  
but the culture was also incubated for about 5 or 9  
hours further at 30C with shaking along with the  
induced cultures. After either 5 or 9 hours  
incubation, the optical density of cultures were  
25 measured at about 600nm, an aliquot of cultures were  
examined by oil emersion microscopy at 1600X  
magnification, and aliquots of cultures were pelleted  
by centrifugation. Bacterial pellets of cultures were  
processed for SDS-polyacrylamide gel electrophoresis on  
30 a 14 percent gel to examine levels of protein produced  
in crude lysates and for N-terminal sequencing  
confirmation of the recombinant gene product. The gel  
was stained with Coomassie blue.

The results are shown in Figure 16, Lanes 8-  
35 10 (human DKR-2 polypeptide) and in Figure 17 (mouse

- 77 -

DKR-2 polypeptide). In Figure 16, Lane 8 contains crude lysate of uninduced control cells; Lane 9 contains crude lysate of induced cells cultured at 30C, and Lane 10 contains crude lysate of induced cells cultured at 37C. The arrow to the left of Lane 10 indicates the expected location of human DKR-2-26-259. As can be seen, significant amounts of polypeptide were generated in the induced cultures whether grown at 30C or 37C, while the uninduced cells did not produce a large amount of polypeptide. Figure 17 shows the results of polypeptide production of mouse DKR-2-26-259. Lane 1 is molecular weight markers. Lanes 2-4 are one clone of *E coli* cells transfected with the DKR-2 plasmid, while Lanes 5-7 are a second clone transfected with the same plasmid. Lanes 2 and 5 are crude lysates of uninduced control cells; Lanes 3 and 6 are crude lysates of induced cells cultured at 30C; and Lanes 4 and 7 are crude lysates of cells cultured at 37C. The arrows to the left of Lanes 4 and 7 indicate the expected location of the DKR-2 polypeptide. As can be seen, large amounts of recombinant protein were observed in crude lysates of induced cultures at 37C but not at 30C. Microscopic analysis of bacterial cells revealed most cells contained at least one inclusion body, suggesting that at least some of the protein may be produced in the insoluble fraction of *E. coli*.

#### Example 9: Expression of DKR-3 in Bacteria

PCR amplification employing the primer pairs and templates described below were used to generate various forms of DKR-3. One primer of each pair introduces a TAA stop codon and a unique *SacII* site following the carboxy terminus of the gene. The other primer of each pair introduces a unique *NdeI* site, a N-

- 78 -

terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, 5 restriction digested, and inserted into the unique NdeI and SacII sites of vector pAMG21 (ATCC accession no. 98113) and transformed into the prototrophic *E. coli* host GM121. Other commonly used *E. coli* expression vectors and host cells are also suitable for expression 10 by one skilled in the art. After transformation, positive clones were selected, plasmid DNA was isolated and the sequence of the DKR-3 gene insert was confirmed.

15 The construct pAMG21-human DKR-3-23-350 was engineered to be 329 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

20 Met-Pro-Ala-Pro-Thr-Ala (SEQ ID NO:55)

Gly-Gly-Glu-Glu-Ile (SEQ ID NO:56).

The template used for PCR was human DKR-3 cDNA and the 25 following oligonucleotides were the primer pair used for PCR and cloning this gene construct.

GTTCTCCTCATATGCCTGCTCCAACTGCAACTTCGGCTCCAGTCAAGCCCCGGCC  
(SEQ ID NO:57)

30 TACGCACTCCGCGGTTAAATCTCTCCCCTCCCAGCA (SEQ ID NO:58)

The construct pAMG21-human DKR-3-33-350 was engineered to be 319 amino acids in length and have the 35 following N-terminal and C-terminal residues, respectively:



Met-Lys-Pro-Gly-Pro-Ala

(SEQ ID NO:59)

Gly-Gly-Glu-Glu-Ile

SEQ ID NO:60)

5

The template used for PCR was human DKR-3 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

10 GTTCTCCTCATATGAAACCAGGTCCAGCCTTAAGCTACCCGCAGGAGGAGGCCA  
(SEQ ID NO:61)

TACGCACTCCGCGGTAAATCTCTTCCCCTCCCAGCA (SEQ ID NO:62)

15

The construct pAMG21-human DKR-3-42-350 was engineered to be 310 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

20 Met-Gln-Glu-Glu-Ala-Thr

(SEQ ID NO:63)

Gly-Gly-Glu-Glu-Ile

(SEQ ID NO:64)

25 The template used for PCR was human DKR-3 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

GTTCTCCTCATATGCAAGAAGAAGCTACTCTGAATGAGATGTTCCGCGAGGTT  
(SEQ ID NO:65)

30

TACGCACTCCGCGGTAAATCTCTTCCCCTCCCAGCA (SEQ ID NO:66)

35 The construct pAMG21-mouse DKR-3-33-349 was engineered to be 318 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

- 80 -

Met-Glu-Pro-Gly-Pro-Ala

(SEQ ID NO:67)

Gly-Glu-Glu-Glu-Ile

(SEQ ID NO:68)

5

The template used for PCR was mouse DKR-3 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

10 GTTCTCCTCATATGGAACCAGGTCCAGCTTTAACTACCCCTCAGGAGGAAGCTA  
(SEQ ID NO:69)

TACGCACTCCGCGGTTAAATCTCCTCCTCTCCGCCTA (SEQ ID NO:70)

15

Transformed *E. coli* GM121 containing the various pAMG21 DKR-3 plasmids described above were grown in 2X YT media containing 20 micrograms/ml kanamycin at 30 °C until the culture reached an optical density at 600 nm of about 0.5. Induction of DKR-3 polypeptide expression was achieved by addition of *Vibrio fischeri* synthetic autoinducer to 100 ng/ml final concentration and incubation of the culture at either 30 or 37C for about 6 hours further with shaking. In addition, as a uninduced control, for each culture no autoinducer was added to an aliquot of the culture, but the culture was also incubated for about 6 hours further at 30C with shaking along with the induced cultures. After about 6 hours, the optical density of cultures were measured at about 600 nm, an aliquot of cultures were examined by oil emersion microscopy at 1600X magnification, and aliquots of cultures were pelleted by centrifugation. Bacterial pellets of cultures were processed for SDS-polyacrylamide gel electrophoresis to examine levels of protein produced in crude lysates, or bacterial pellets were processed to determine whether the recombinant

20

25

30

35

- 81 -

protein was in the soluble or insoluble fraction of *E. coli* and for N-terminal sequencing confirmation of the recombinant gene product. The results are shown as photos of the SDS gels in Figures 18 and 19. In Figure 18, Lane 10 is molecular weight markers, and Lanes 1-9 are crude lysates of bacterial cells. Lane 1 is crude lysate of uninduced control cells; Lanes 2, 4, 6, and 8 are crude lysates of induced cells cultured at 30C; Lanes 3, 5, 7, and 9 are induced cells cultured at 37C. Lanes 1-5 contain lysates of cells transfected with the pAMG21-human DKR-3-23-350 construct; and Lanes 6-9 contain lysates of cells transfected with the pAMG21-human DKR-3-33-350 construct. The arrows to the left of Lane 2 and the right of Lane 9 indicate the expected location of the DKR-3 polypeptides. Figure 19 contains molecular weight markers in Lane 10; Lanes 1-5 are crude lysates of cultured cells transfected with the pAMG21-human DKR-3-42-350 construct; Lanes 6-9 are crude lysates of cells transfected with the pAMG21-mouse DKR-3-33-349 construct. Lanes 1 and 6 are uninduced controls; Lanes 2, 4, 7, and 8 are crude lysates of induced cells cultured at 30C (two different clones of each construct); Lanes 3, 5, and 9 are crude lysates of induced cells cultured at 37C (two separate clones of the human DKR-3-42-350 construct in Lanes 3 and 5). The arrow to the right of Lane 9 indicates the expected location of the mouse DKR-3 polypeptides; the arrow to the left of Lane 4 indicates the expected location of human DKR-3 polypeptide. As can be seen, all DKR-3 constructs produced large amounts of recombinant protein in *E. coli*. No inclusion bodies could be detected by oil emersion microscopy, and the recombinant polypeptides were mostly found in the soluble fraction of the cells.

- 82 -

Example 10: Expression of DKR-4 in Bacteria

PCR amplification employing the primer pairs and template described below were used to generate a recombinant form of human DKR-4. One primer of each pair introduces a TAA stop codon and a unique *Bam*HI site following the carboxy terminus of the gene. The other primer of each pair introduces a unique *Nde*I site, a N-terminal methionine, and optimized codons for the amino terminal portion of the gene. PCR and thermocycling was performed using standard recombinant DNA methodology. The PCR products were purified, restriction digested, and inserted into the unique *Nde*I and *Bam*HI sites of the vector pAMG21 (ATCC accession no. 98113) and transformed into the prototrophic *E. coli* host GM94. Other commonly used *E. coli* expression vectors and host cells are also suitable for expression. After transformation, positive clones were selected and will be examined for expression of the recombinant gene product.

The construct pAMG21-human DKR-4-19-224 was engineered to be 207 amino acids in length and have the following N-terminal and C-terminal residues, respectively:

Met-Leu-Val-Leu-Asp-Phe

(SEQ ID NO:71)

Lys-Ile-Glu-Lys-Leu

(SEQ ID NO:72)

The template used for PCR was human DKR-4 cDNA and the following oligonucleotides were the primer pair used for PCR and cloning this gene construct:

- 83 -

GTTCTCCTCATATGTTAGTTTTGGATTTCAACAACATCAGGAGCTCT  
(SEQ ID NO:73)

TACGCACTGGATCCTTACAGTTTTTCTATTTTTTGGCATACTCTTAATC  
5 (SEQ ID NO:74)

It is anticipated that DKR-4 polypeptide could be prepared using the PCR product as described above for the other DKR polypeptides.

10

EXAMPLE 11: Production and Purification of DKR-3 Polypeptide in Mammalian Cells

Human DKR-3 cDNA was cloned onto the  
15 mammalian expression vector pcDNA3.1(-)/mycHis (Invitrogen, Carlsbad, CA) and the vector construct was amplified using the Qiagen maxi-prep kit (Qiagen, Chatsworth, CA) standard ligation techniques.

Human embryonic kidney 293T cells (American  
20 Type Culture Collection) were cultured in 10 cm dishes, and grown to about 80 percent confluence. The cells were then transfected with the vector construct using the DMRIE-C® liposome formulation (Gibco BRL, Grand Island, NY) as follows. About 240 microliters of  
25 DMRIE-C® were added to 8 ml of Optimem medium. About 40 ul (equivalent to about 56 micrograms) of purified vector construct was then added to another 8 ml of Optimem. After mixing and incubation at room temperature for about 15 minutes, 2 ml of this solution  
30 was added to each of 8 plates. After about 5 hours, the medium was aspirated and 10 ml of DME medium containing about 10 percent fetal calf serum was added. The cells were incubated 16-18 hours after which the medium was removed and about 10 ml of SF Optimem medium  
35 per well without phenol red were added. After about 24 hours, this medium, the "conditioned medium" was

- 84 -

harvested, passed over a 0.22 micron filter and stored at 4°C. The cells were then incubated in another 10 ml of SF Optimem per plate. After 24 hours, this medium was collected, filtered and also stored at 4°C.

5           The conditioned media was added to a buffer containing 50 mM NaPO<sub>4</sub>, pH8, and 250 mM sodium chloride, and passed over a column of nickel-Sephadex (Qiagen, Chatsworth, CA). Non-specifically bound proteins were eluted using the same buffer containing  
10 10 mM imidazole, followed by the same buffer containing 20 mM imidazole. DKR-3 was then eluted using 125 mM-250 mM imidazole. Fractions from the column were subjected to 12 percent SDS gel electrophores and silver stained. The results are shown in Figure 20.  
15 Lane 2 contains material that was loaded on to the gel. Lane 3 contains the flow through fraction after loading the column with conditioned medium, Lanes 4, 5, 6, and 7 contain column fractions after treatment with 10, 20, 125, and 250 mM imidazole. Molecular weight standards  
20 are shown in Lane 8. As can be seen a single band of protein of the correct molecular weight is seen in Lanes 5 and 6, indicating that this procedure resulted in generation of purified DKR-3 protein (attached to myc and His tags). Smearing of the protein band may be  
25 due to glycosylation. Separately, a Western blot was run to confirm that the purified protein did indeed have a His tag (indicating that the fusion protein DKR-3 mycHis had been produced). The Western blot was prepared using standard procedures and was probed with  
30 a polyclonal anti-His-HRP antibody (Invitrogen, Carlsbad, CA). A photo of the Western blot is shown in Figure 21; the Lanes correspond to that for the gel (described immediately above). As can be seen, there is antibody binding in Lanes 2, 5, and 6, indicating  
35 that DKR-3 mycHis was loaded on to the column and was eluted in the 20 and 125 mM imidazole washes.

- 85 -

Example 12: Anchorage Independent Growth Assay

5 A distinguishing feature of many cancer cell  
lines is their ability to grow in an anchorage  
independent manner. Whereas normal cells will only  
grow and divide until they come in contact with their  
neighbors, cancer cells continue to grow and divide  
after contact, thereby forming tumors. Thus, one assay  
10 for cancer cell growth inhibitor compounds measures the  
ability of cancer cells to grow and divide in the  
presence of the compound. There are many ways known to  
the skilled artisan in which this assay can be  
conducted, however two preferred methods are set forth  
15 below.

A. Stably Transfected Cell Assay

In this procedure, any human cancer cell line  
that does not express the DKR gene to be tested (either  
20 human DKR-1, 2, 3, 4, or a fragment or variant thereof)  
is transfected with the DKR gene under evaluation,  
where the DKR gene is inserted into a vector such as  
pcDNA3.1 (Invitrogen, Carlsbad, CA) or other suitable  
mammalian expression vector. Transfection can be  
25 conducted as described herein. The transfected cancer  
cells are cultured to generate a stably transfected  
cell line. Once a stably transfected cell line has  
been established, the cells are added to Noble or  
equivalent agar (about 0.35 percent) prepared in  
30 standard mammalian cell culture medium such as RPMI.  
The cell/agar solution is poured on to petri plates  
containing solidified agar ban (about 0.5 percent  
agar). Colony formation is evaluated daily to  
determine the rate of growth of the cells, and culture  
35 medium is added to each plate as needed. Separately,  
the same cells are transfected with vector only

- 86 -

(containing no DKR gene). These "control" cells are then treated in an identical manner to the DKR gene containing cells and can be used as a standard of comparison for the DKR gene containing cells.

- 5           Examples of suitable cancer cell lines for conducting this assay include, without limitation, human breast cancer cell line MCF7 and the glioblastoma cell line U-87MG.

10   B. Protein Assay

- An alternate or additional assay to measure the growth of cancer cell lines treated with a DKR polypeptide is as follows. Any human cancer cell line  
15   not expressing the DKR polypeptide under evaluation can be cultured and prepared with an agar solution as described above. The cells can then be plated as described, and a solution of DKR polypeptide (either full length, or a fragment or variant thereof) in  
20   culture medium can be added to the agar either daily, every other day, or once per week for three weeks. Typically, a concentration of about 10 nM will be added, although a series of dilutions ranging from 1 nM to 1 mM can be used. Control plates will receive a  
25   solution of culture medium only. The plates can be monitored daily for up to about three weeks to evaluate cell colony formation. After three weeks, control and experimental plates can be compared for the number and size of cell colonies. It is anticipated that those  
30   plates receiving DKR polypeptide that is biologically active will have fewer cell colonies, and the colonies will be smaller, as compared to control plates.



- 87 -

Example 13: In Vivo Tumor Assay

The ability of each DKR polypeptide to inhibit tumor growth *in vivo* can be evaluated as follows. Tumor cells not expressing the DKR gene under evaluation can be transfected using procedures described herein with a DKR nucleic acid construct encoding a full length DKR gene, or a fragment or variant thereof. The transfected cells can be maintained in culture (as described herein) until ready for use.

Male or female athymic nude mice (Charles River Labs, Boston, MA) are kept in a sterile environment. The mice are then injected with about  $2 \times 10^6$  cells (either DKR transfected cells or control "vector only" transfected cells) in a total volume of about 0.1 ml can be injected sub-cutaneously. The mice can then be examined daily for appearance of (a) tumor(s) and for the size of the tumor. Preferably, the mice will be examined for up to about six months so as to provide time for tumor growth (and regression where DKR polypeptides are effective at decreasing tumor growth). The tumor(s), where present, can then be removed, weighed and examined for (1) the presence of DKR polypeptide, and (2) morphology. Tumors from mice containing DKR construct transfected cells can be compared to tumors from mice containing cells transfected with vector only. It is anticipated that DKR polypeptides, due to their similarity with *dkk-1*, a potent *wnt8* antagonist, will be able to decrease the size of the tumor as compared with controls.

- 88 -

We Claim:

1. An isolated nucleic acid molecule  
encoding a biologically active DKR polypeptide selected  
5 from the group consisting of:
  - (a) the nucleic acid molecule comprising SEQ  
ID NO:1;
  - (b) the nucleic acid molecule comprising SEQ  
ID NO:2;
  - 10 (c) the nucleic acid molecule comprising SEQ  
ID NO:3;
  - (d) the nucleic acid molecule comprising SEQ  
ID NO:4;
  - (e) the nucleic acid molecule comprising SEQ  
15 ID NO:5;
  - (f) the nucleic acid molecule comprising SEQ  
ID NO:6;
  - (g) the nucleic acid molecule comprising SEQ  
ID NO:7;
  - 20 (h) the nucleic acid molecule comprising SEQ  
ID NO:75;
  - (i) the nucleic acid molecule comprising SEQ  
25 ID NO:76;
  - (j) the nucleic acid molecule comprising SEQ  
ID NO:77;
  - 30 (k) the nucleic acid molecule comprising SEQ  
ID NO:78;
- (l) the nucleic acid molecule encoding the  
polypeptide of SEQ ID NO:8;
- 35 (m) a nucleic acid molecule encoding the  
polypeptide of SEQ ID NO:9;

- 89 -

(n) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:10, or a biologically active fragment thereof;

5 (o) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:11, or a biologically active fragment thereof;

(p) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:12, or a biologically active fragment thereof;

10 (q) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:13, or a biologically active fragment thereof;

(r) a nucleic acid molecule encoding the polypeptide of SEQ ID NO:14, or a biologically active  
15 fragment thereof

(s) a nucleic acid molecule that encodes a polypeptide that is at least 85 percent identical to the polypeptide of SEQ ID NOS: 10, 11, 12, 13, or 14;

(t) a nucleic acid molecule that encodes a  
20 biologically active DKR polypeptide that has 1-100 amino acid substitutions and/or deletions as compared with the polypeptide of any of SEQ ID NOS:8, 9, 10, 11, 12, 13, or 14; and

(u) a nucleic acid molecule that hybridizes  
25 under conditions of high stringency to any of (c), (d), (e), (f), (g), (h), (i), (k), (l), (m), (n), (o), (p), (q), (r), (s), and (t) above.

2 An isolated nucleic acid molecule that is  
30 the complement of the nucleic acid molecule of claim 1.

3. An isolated nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or SEQ ID NO:7.

35

- 90 -

4. An isolated nucleic acid molecule encoding the polypeptide of SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, or SEQ ID NO:14.

5

5. An isolated nucleic acid molecule encoding a biologically active DKR polypeptide selected from the group consisting of: amino acids 16-350, 21-350, 22-350, 23-350, 33-350, or 42-350, 21-145, 40-145, 40-150, 45-145, 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, or 310-350 of SEQ ID NO:9; amino acids 15-266, 24-266, or 32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or 34-359 of SEQ ID NO:12; and amino acids 19-224, 20-224, 21-224, or 22-224 of SEQ ID NO:14.

15

6. A vector comprising the nucleic acid molecule of claim 1.

7. A vector comprising the nucleic acid molecule of claim 2.

20

8. A vector comprising the nucleic acid molecule of claim 3.

9. A vector comprising the nucleic acid molecule of claim 4.

25

10. A vector comprising the nucleic acid molecule of claim 5.

30

11. A host cell comprising the vector of claim 6.

12. A host cell comprising the vector of claim 7.

35

- 91 -

13. A host cell comprising the vector of  
claim 8.

14. A host cell comprising the vector of  
5 claim 9.

15. A host cell comprising the vector of  
claim 10.

16. A process for producing a biologically  
active DKR polypeptide comprising the steps of:  
    (a) expressing a polypeptide encoded by the  
nucleic acid of claim 1 in a suitable host; and  
    (b) isolating the polypeptide.

17. The process of claim 16 wherein the  
polypeptide is SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10,  
SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13 or SEQ ID  
NO:14.

18. A biologically active DKR polypeptide  
selected from the group consisting of:  
    (a) the polypeptide of SEQ ID NO:8;  
    (b) the polypeptide of SEQ ID NO:9;  
    (c) the polypeptide of SEQ ID NO:10;  
    (d) the polypeptide of SEQ ID NO:11;  
    (e) the polypeptide of SEQ ID NO:12;  
    (f) the polypeptide of SEQ ID NO:13;  
    (g) the polypeptide of SEQ ID NO:14;  
    (h) a polypeptide that has 1-100 amino acid  
substitutions or deletions as compared with the  
polypeptide of any of (a)-(h) above; and  
    (i) a polypeptide that is at least 85 percent  
identical to any of the polypeptides of (c)-(h) above.

19. The polypeptide of claim 18 that does  
not possess an endogenous signal peptide.

- 92 -

20. A polypeptide selected from the group  
consisting of amino acids 16-350, 21-350, 22-350, 23-  
350, 33-350, 42-350, 21-145, 40-145, 40-150, 45-145,  
5 45-145, 145-290, 145-300, 145-350, 150-290, 300-350, or  
310-350 of SEQ ID NO:9; amino acids 15-266, 24-266, or  
32-266 of SEQ ID NO:10; amino acids 17-259, 26-259, or  
34-259 of SEQ ID NO:12; and amino acids 19-224, 20-224,  
21-224, or 22-224 of SEQ ID NO:14.

10

1 / 25

FIG. 1

1 ATGCAGCGC TCGGGGGTAT TTTGCTGTGT AACTGCTGG CGCGGGCGGT  
51 CCCCACTGCT CCTGCTCCTT CCCGACGGT CACTTGGACT CCGGGGGAGC  
101 CGGGCCAGC TCTCAACTAC CCTCAGGAGG AAGCTACGCT CAATGAGATG  
151 TTTCGAGAGG TGGAGGAGCT GATGGAAGC AGCTGCTAAA AACTGCGCAG  
201 TGCCGTGGAG GAGATGGAGG CGGAAGAAGC AGCTGCTAAA AACTGCGCAG  
251 AGGTGAACCT GGCAAGCTTA CCTCCAACT ATCACAATGA GACCAGCAGC  
301 GAGACCAGGG TGGGAATAA CACAGTCCAT GTGCACCAGG AAGTTCACAA  
351 GATAACCAAC AACCAGAGTG GACAGGTGGT CTTTCTTGAG ACAGTCATTA  
401 CATCTGTAGG GGATGAAGAA GGCAAGAGGA GCCATGAATG TATCATTTGAT  
451 GAAGACTGTG GGCCACCCAG GACTGCCAG TTCTCCAGCT TCAAGTACAC  
501 CTGCCAGCCA TGCCGGGACC AGCAGATGCT ATGCACCCGA GACAGTGAGT  
551 GCTGTGGAGA CCAGCTGTGT GCCTGGGGTC ACTGCACCCA AAAGGCCACC  
601 AAAGGTGGCA ATGGGACCAT CTGTGACAA CAGAGGGATT GCCAGCCTGG  
651 CCTGTGTTGT GCCTTCCAAA GAGGCCCTGCT GTTCCCGGTG TGCACACCCC  
701 TGCCCGTGGA GGGAGAGCTC TGCCATGACC CCACCAGCCA GCTGCTGGAT  
751 CTCATCACCT GGGAACTGGA GCCTGAAGGA GCTTTGGACC GATGCCCTTG  
801 CGCCAGTGGC CTCCTATGCC AGCCACACAG CCACAGTCTG GTGTACATGT  
851 GCAAGCCAGC CTTGCTGGG AGCCATGACC ACAGTGAGGA GAGCCAGCTG  
901 CCCAGGGAGG CCCCGGATGA GTACGAAGAT GTTGGCTTCA TAGGGGAAGT  
951 GCGCCAGGAG CTGGAAGACC TGGAGCGGAG CCTAGCCCCAG GAGATGGCAT  
1001 TTGAGGGGCC TGCCCCCTGTG GAGTCACTAG GCGGAGAGGA GGAGATTAG

FIG. 2

```

1  ATGCAGCGGC TTGGGGCCAC CCTGCTGTGC CTGCTGCTGG CGGCGGCGGT
51  CCCACGGCC CCGCGGCCG CTCCGACGGC GACCTCGGCT CCAGTCAAGC
101 CCGGCCCCGGC TCTCAGCTAC CCGCAGGAGG AGGCCACCCT CAATGAGATG
151 TTCCGCGAGG TTGAGGAACT GATGGAGGAC ACGCAGCACA AATTGCGCAG
201 CGCGGTGGAA GAGATGGAGG CAGAAGAAGC TGCTGCTAAA GCATCATCAG
251 AAGTGAACCT GGCAAACTTA CCTCCCAGCT ATCACAATGA GACCAACACA
301 GACACGAAGG TTGGAATAA TACCATCCAT GTGCACCGAG AAATTCACAA
351 GATAACCAAC AACCAGACTG GACAAATGGT CTTTTCAGAG ACAGTTATCA
401 CATCTGTGGG AGACGAAGAA GGCAGAAGGA GCCACGAGTG CATCATCGAC
451 GAGGACTGTG GGCCAGCAT GTACTGCCAG TTTGCCAGCT TCCAGTACAC
501 CTGCCAGCCA TGCCGGGGCC AGAGGATGCT CTGCACCCGG GACAGTGAGT
551 GCTGTGGAGA CCAGCTGTGT GTCTGGGGTC ACTGCACCAA AATGGCCACC
601 AGGGGCAGCA ATGGGACCAT CTGTGACAA CAGAGGGAAT GCCAGCCGGG
651 GCTGTGCTGT GCCTTCCAGA GAGGCCTGCT GTTCCCTGTG TGCACACCCC
701 TGCCCGTGGA GGGCGAGCTT TGCCATGACC CCGCCAGCCG GCTTCTGGAC
751 CTCATCACCT GGGAGCTAGA GCCTGATGGA GCCTTGACC GATGCCCTTG
801 TGCCAGTGGC CTCCCTCTGCC AGCCCCACAG CCACAGCCTG GTGTATGTGT
851 GCAAGCCGAC CTTCGTGGG AGCCGTGACC AAGATGGGA GATCCTGCTG
901 CCCAGAGAGG TCCCCGATGA GTATGAAGTT GGCAGCTTCA TGGAGGAGGT
951 GCGCCAGGAG CTGGAGGACC TGGAGAGGAG CCTGACTGAA GAGATGGCGC
1001 TGGGGGAGCC TGCGGCTGCC GCCGCTGCAC TGCTGGGAGG GGAAGAGATT
1051 TAG

```



3 / 25

FIG. 3

```

1  ATGATGGCTC TGGGCGCAGC GGGAGCTACC CGGGTCTTTG TCGCGATGGT
51  AGCGGCGGCT CTGGGCGGCC ACCCTCTGCT GGGAGTGAGC GCCACCTTGA
101  ACTCGGTTCT CAATTCCAAC GCTATCAAGA ACCTGCCCCC ACCGCTGGGC
151  GCGCTGCGG  GGCACCCAGG CTCTGCAGTC AGCGCCGCGC CGGGAATCCT
201  GTACCCGGGC  GGAATAAGT ACCAGACCAT TGACAACCTAC CAGCCGTACC
251  CGTGCGCAGA GGACGAGGAG TCGGGCACTG ATGAGTACTG CGCTAGTCCC
301  ACCCGCGGAG GGGACGCGGG CGTGCAAATC TGTCTCGCCT GCAGGAAGCG
351  CCGAAAACGC TGCATGCCGC ACGCTATGTG CTGCCCCGGG AATTACTGCA
401  AAAATGGAAT ATGTGTGTCT TCTGATCAAA ATCATTTCCG AGGAGAAATT
451  GAGGAAACCA TCACTGAAAG CTTTGGTAAT GATCATAGCA CCTTGGATGG
501  GTATTCCAGA AGAACCACCT TGTCTTCAA AATGTATCAC ACCAAAGGAC
551  AAGAAGGTTT TGTGTGTCTC CGGTCATCAG ACTGTGCCTC AGGATTGTGT
601  TGTGCTAGAC ACTTCTGGTC CAAGATCTGT AAACCTGTCC TGAAGAAGG
651  TCAAGTGTGT ACCAAGCATA GGAGAAAAGG CTCTCATGGA CTAGAAATAT
701  TCCAGCGTTG TTAAGTGTGA GAAGGTCTGT CTTGCCGGAT ACAGAAAGAT
751  CACCATCAAG CCAGTAATTC TTCTAGGCTT CACACTTGTC AGAGACACTA
801  A

```

FIG. 4

1 ATGGCCGGC TGATCGGGT CAAGGATCA TCCGCTGCC TTCTCCTACT  
51 GGCCGGGTG CTGATGGTGG AGAGCTACA GCTAGGCAGC TCGCGGGCCA  
101 AACTCAACTC CATCAAGTCC TCTCTAGGAG GGGAGACTCC TGCTCAGTCA  
151 GCCAACCGAT CTGCAGGCAT GAACCAAGGA CTGGCTTTCG GCGGCAGTAA  
201 GAAGGGCAA AGCCTGGGC AGGCCTACCC TTGCAGCAGT GATAAGGAAT  
251 GTGAAGTTGG AAGATACTGC CACAGTCCCC ACCAAGGATC ATCAGCCTGC  
301 ATGCTCTGTA GGAGGAAAA GAAACGATGC CACAGAGATG GGATGTGTG  
351 CCTTGGTACC CGCTGCAATA ATGGAATCTG CATCCCAGTC ACTGAGAGCA  
401 TCCTCACCCC ACATATCCCA GCTCTGGATG GCACCCGGCA TAGAGATCGC  
451 AACCATGGTC ACTATTCCAA CCATGACCTG GGATGGCAGA ATCTAGGAAG  
501 GCCACACTCC AAGATGCCCTC ATATAAAGG ACATGAAGGA GACCCATGCC  
551 TACGGTCATC AGACTGCATT GATGGGTTTT GTTGTGCTCG CCACTTCTGG  
601 ACCAAATCT GCAAACCCAGT GCTCCATCAG GGGGAAGTCT GTACCAACA  
651 ACGCAAGAAG GGTTCGCACG GGCTGGAGAT TTTCAGAGG TGTGACTGTG  
701 CAAAGGCCT GTCCCTGCAA GTGTGGAAG ATGCCACCTA CTCTTCCAAA  
751 GCCAGACTCC ATGTATGCCA GAAGATCTGA

FIG. 5

1 ATGGCCGCGT TGATCGGGAG CAAGGATTGG TCCTGCTGCC TGCTCCTACT  
51 GGCCGCGGTG CTGATGGTGG AGAGCTCACA GATCGGCAGT TCGCGGGCCA  
101 AACTCAACTC CATCAAGTCC TCTCTGGCG GGGAGACGCC TGGTCAGGCC  
151 GCCAATCGAT CTGCGGGCAT GTACCAAGGA CTGGCATTGG GCGGCAGTAA  
201 GAAAGGCAAA AACCTGGGSC AGGCCCTACCC TTGTAGCAGT GATAAGGAGT  
251 GTGAAGTTGG GAGGTATTGC CACAGTCCCC ACCAAGGATC ATCGGCCTGC  
301 ATGGTGTGTC GGAGAAAAA GAAGCGCTGC CACCGAGATG GCATGTGCTG  
351 CCCCAGTACC CGCTGCAATA ATGGCATCTG TATCCCAGTT ACTGAAAGCA  
401 TCTTAACCCC TCACATCCC GCTCTGGATG GTCCTCGCA CAGAGATCGA  
451 AACCACGGTC ATTACTCAA CCATGACTTG GGATGGCAGA ATCTAGGAAG  
501 ACCACACACT AAGATGTCAC ATATAAAGG GCATGAAGGA GACCCCTGCC  
551 TACGATCATC AGACTGCATT GAAGGGTTTT GCTGTGCTCG TCATTCTGG  
601 ACCAAAATCT GCAAACCAAGT GCTCCATCAG GGGGAAGTCT GTACCAACA  
651 ACGCAAGAAG GGTCTCATG GGCTGGAAAT TTTCCAGCGT TGCAGCTGTG  
701 CGAAGGGCCT GTCTTGCAA GTATGGAAG ATGCCACCTA CTCCTCCAAA  
751 GCCAGACTCC ATGTGTGTCA GAAAATTGA

6 / 25

FIG. 6

```
1  ATGGCCGCGT TGATGCGGAG CAAGGATTGG TCCTGCTGCC TGCTCCTACT
51  GGCCGCGGTG CTGATGGTGG AGAGCTCACA GATCGGCAGT TCGGGGGCCA
101  AACTCAACTC CATCAAGTCC TCTCTGGGCG GGGAGACGCC TGGTCAGGCC
151  GCCAATCGAT CTGCGGGCAT GTACCAAGGA CTGGCATTCG GCGGCAGTAA
201  GAAGGGCAAA AACCTGGGGC AGCCTACCC TTGTAGCAGT GATAAGGAGT
251  GTGAAGTTGG GAGGTATTGC CACAGTCCC ACCAAGGATC ATCGGCCCTGC
301  ATGGTGTGTC GGAGAAAAAA GAAGCGCTGC CACCGAGATG GCATGTGCTG
351  CCCCAGTACC CGCTGCAATA ATGGGCATGA AGGAGACCCC TGCCCTACGAT
401  CATCAGACTG CATTGAAGGG TTTTGCTGTG CTCGTCATTT CTGGACCAAA
451  ATCTGCAAAC CAGTGCTCCA TCAGGGGGAA GTCTGTACCA AACAAACGCAA
501  GAAGGGTTCT CATGGGCTGG AAATTTTCCA GCGTTGCGAC TGTGCGAAGG
551  GCCTGTCTTG CAAAGTATGG AAAGATGCCA CCTACTCCTC CAAAGCCAGA
601  CTCCATGTGT GTCAGAAAAT TTGA
```

FIG. 7

```

1  ATGGTGGCGG CCGTCCCTGCT GGGGCTGAGC TGGCTCTGCT CTCCCCTGGG
51  AGCTCTGGTC CTGGACTTCA ACAACATCAG GAGCTCTGCT GACCTGCATG
101  GGGCCCCGGAA GGGCTCACAG TGCCTGTCTG ACACGACTG CAATACCAGA
151  AAGTTCTGCC TCCAGCCCCG CGATGAGAAG CCGTCTCTGT CTACATGTCTG
201  TGGGTTGCGG AGGAGGTGCC AGCGAGATGC CATGTGCTGC CCTGGGACAC
251  TCTGTGTGAA CGATGTTTGT ACTACGATGG AAGATGCAAC CCCAATATTA
301  GAAAGGCAGC TTGATGAGCA AGATGSCACA CATGCAGAAG GAACAACTGG
351  GCACCCAGTC CAGGAAAACC AACCCAAAAG GAAGCCAAGT ATTAAGAAAT
401  CACAAGGCAG GAAGGGACAA GAGGGAGAAA GTTGTCTGAG AACTTTTGAC
451  TGTGGCCCTG GACTTTGCTG TGCTCGTCAT TTTTGGACGA AAATTTGTAA
501  GCCAGTCCTT TTGGAGGGAC AGGTCTGCTC CAGAAGAGGG CATAAAGACA
551  CTGCTCAAGC TCCAGAAATC TTCCAGCGTT GCGACTGTGG CCCTGGACTA
601  CTGTGTCGAA GCCAATTGAC CAGCAATCGG CAGCATGCTC GATTAAAGAGT
651  ATGCCAAAAA ATAGAAAAGC TATAA

```

## FIG. 8

1 MORLGGILLC TLLAAAVPTA PAPSPTVTWT PAEPGPALNY PQEEATLNEM  
51 FREVEELMED TQHKLRSAVE EMEAEAAAK TSSEVNLA SL PPNYHNETST  
101 ETRVGNNTVH VHQEVHKITN NQSGQVVFSE TVITSVGDEE GKRSHECIID  
151 EDCGPTRYCQ FSSFYTCQP CRDQQLCTR DSECCGDQLC AWGHCTQKAT  
201 KGGNGTICDN QRDCCPGLCC AFQGLLFPV CTPLPVEGEL CHDPTSQLLD  
251 LITWELEPEG ALDRCPGASG LLCQPHSHSL VYMCKPAFVG SHDHSEESQL  
301 PREAPDEYED VGFIGEVRQE LEDLERSLAQ EMAFEGPAPV ESLGGEETI\*

9 / 25

## FIG. 9

1 MORLGATLLC LLLAAAVPTA PAPAPTATSA PVKPGPALSY PQEEATLNEM  
51 FREVEELMED TQHKLRSAVE EMEAEAAAK ASSEVNLANL PPSYHNETNT  
101 DTKVGNNTH VHREIHKITN NQTGQMFSE TVITSVGDEE GRRSHECIID  
151 EDCGPSMYCQ FASFQYTCQP CRGQRMCLCTR DSECCGDQLC VWGHCTKMAT  
201 RGSNGTICDN QRDCQPLCC AFQGLLFPV CTPLPVEGEL CHDPASRLLD  
251 LITWELEPDG ALDRCPGASG LLCQPHSHSL VYVCKPTFVG SRDQDGEILL  
301 PREVPDEYEV GSFMEEVQRQE LEDLERSLTE EMALGEPAAA AAALLGEEI  
351 \*

10 / 25

## FIG. 10

1	MMALGAAGAT	RVEVAMVAAA	LGGHPLLGV	ATLNSVLNSN	AIKNLPPPLG
51	GAAGHPGSAV	SAAPGILYPG	GNKYQTIDNY	QYPYCAEDEE	CGTDEYCASP
101	TRGGDAGVQI	CLACRKRKR	CMRHAMCCPG	NYCKNGICVS	SDQNHFRGEI
151	EETITESFGN	DHSTLDGYSR	RTTLSSKMYH	TKQEGSVCL	RSSDCASGLC
201	CARHFWSKIC	KPVLKEGQVC	TKHRRKGSHG	LEIFQRCYCG	EGLSCRIQKD
251	HHQASNSSRL	HTCQRH*			



11 / 25

## FIG. 11

1	MAALMRVKDS	SRCLLLLAAV	LMVSSQLGS	SRAKLNSIKS	SLGGETPAQS
51	ANRSAGMNQG	LAFGSKKKG	SLGQAYPCSS	DKECEVGRYC	HSPHQGSSAC
101	MLCRRKKKRC	HRDGMCCPGT	RCNNGICIPV	TESILTPHIP	ALDGTRHRDR
151	NHGHYSNHD	L GWQNLGRPHS	KMPHIKGHEG	DPCLRSSDCI	DGFCCARHFW
201	TKICKPVLHQ	GEVCTKQKK	GSHGLEIFQR	CDCAKGLSCK	VWKDATYSSK
251	ARLHVCQKI*				

12 / 25

## FIG. 12

1	MAALMRSKDS	SCCLLLAAV	LMVSSQIGS	SRAKLNSIKS	SLGGETPGQA
51	ANRSAGMYQG	LAFGSKKKG	NLGQAYPCSS	DKECEVGRYC	HSPHQGSSAC
101	MVCRKKKRC	HRDGMCCPST	RCNNGICIPV	TESILTSHIP	ALDGTRHRDR
151	NHGHYSNHD	GWQNLGRPHT	KMSHIKGHEG	DPCLRSSDCI	EGFCCARHEW
201	TKICKPVLHQ	GEVCTQRKK	GSHGLEIFQR	CDCAKGLSCK	VWKDATYSSK
251	ARLHVCQKI*				

13 / 25

## FIG. 13

1 MAALMRSKDS SCCLLLAAV IMVSSQIGS SRAKLNSIKS SLGGETPGQA  
51 ANRSAGMYQG LAFGGSKKGK NLGQAYPCSS DKECEVGRYC HSPHQGSSAC  
101 MVCRRKKKRC HRDGMCCPST RCNNGHEGDP CLRSSDCIEG FCCARHFWTK  
151 ICKPVLHQGE VCTKQKKKGS HGLEIFQRCD CAKGLSCKVW KDATYSSKAR  
201 LHVCQKI\*

1 4 / 2 5

## FIG. 14

1 MVAAVLLGLS WLCSPGLGALV LDFNNIRSSA DLHGARKGSQ CLSDTDCNTR  
51 KFCLQPRDEK PFCATCRGLR RRCQORDAMCC PGTLCVNDVC TTMEDATPIL  
101 ERQLDEQDGT HAEGTTGHPV QENQPKRKPS IKKSQGRKGQ EGESCLRTFD  
151 CGPGLCCARH FWTKICKPVL LEGQVCSRRG HKDTAQAPEI FQRCDCGPGGL  
201 LCRSQTLSNR QHARLRVCQK IEKL\*

FIG. 15A




FIG. 15B




FIG. 15D

FIG. 15C



FIG. 16




FIG. 17




FIG. 18






FIG. 19




FIG. 20





FIG. 21



Imidazole

22 / 25

FIG. 22

1 ATGATGGCTC TGGGTGCTGC TGGTGCTACC CGTGTTTTCG TTGCTATGGT  
 51 TGCTGCTGCT CTGGGTGGTC ACCCGCTGCT GGGTGTTTCC GCTACCCCTGA  
 101 ACTCCGTTCT GAACTCCAAC GCTATCAAAA ACCTGCCGCC GCCGCTGGGT  
 151 GGTGCTGCTG GTCACCCGGG TTCCGCTGTT TCCGCTGCTC CGGGTATCCT  
 201 GTACCCGGGT GGTAAACAAAT ACCAGACCAT CGACAACTAC CAGCCGTACC  
 251 CGTGCGCTGA AGACGAAGAA TCGGGTACCG ACGAATACTG CGCTTCCCCG  
 301 ACCCGTGGTG GTGACGCTGG TGTTCAGATC TGCCTGGCTT GCCGTAAACG  
 351 TCGTAAACGT TGCATGCCGC ACGCTATGTG CTGCCCGGGT AACTACTGCA  
 401 AAAACGGTAT CTGCGTTTCC TCCGACCAGA ACCACTTCCG TGGTGAAATC  
 451 GAAGAAACCA TCACCGAATC CTTGGGTAAC GACCACTCCA CCTGGACGG  
 501 TTAATCCCCG CTACCAACCC TGTCCCTCCAA AATGTACCAC ACCAAAGGTC  
 551 AGGAGGTTT CATTGTCCTG CGTTCCTCCG ACTGCGCTTC CGGTCTGTGC  
 601 TGCCTCGTC ACTTCTGGTC CAAAATCTGC AAACCGGTTT TGAAAGAAGG  
 651 TCAGGTTTGC ACCAAACACC GTCGTAAAGG TTCCCACGGT CTGGAATCT  
 701 TCCAGCGTTG CTACTGCGGT GAAGGTCTGT CCGCCGTAT CCAGAAAAGAC  
 751 CACCACCAGG CTTCCAACTC CTCCCGTCTG CACACCTGCC AGCGTCAC

23 / 25

## FIG. 23

1 ATGGCTGCTC TGATGCGTTC CAAAGACTCC TCCTGCTGCC TGCTGCTGCT  
51 GGCTGCTGTT CTGATGGTTG AATCCTCCCA GATCGGTTCC TCCCGTGCTA  
101 AACTGAACTC CATCAAATCC TCCCTGGGTG GTGAAACCCC GGGTCAGGCT  
151 GCTAACCGTT CCGCTGGTAT GTACCAGGGT CTGGCTTTCG GTGGTTCCAA  
201 AAAAGGTAAA AACCTGGTC AGGCTTACCC GTGCTCCTCC GACAAAGAAT  
251 GCGAAGTTGG TCGTTACTGC CACTCCCCGC ACCAGGGTTC CTCCGCTTGC  
301 ATGGTTTGCC GTCGTAAAAA AAAACGTTGC CACCGTGACG GTATGTGCTG  
351 CCCGTCCACC CGTTGCAACA ACGGTATCTG CATCCCGGTT ACCGAATCCA  
401 TCCTGACCCC GCACATCCCG GCTCTGGACG GTACCCCGTCA CCGTGACCGT  
451 AACCACGGTC ACTACTCCAA CCACGACCTG GGTGGCAGA ACCTGGGTG  
501 TCCGCACACC AAAATGTCCC ACATCAAAGG TCACGAAAGT GACCCGTGCC  
551 TCGGTTCCCTC CGACTGCATC GAAGGTTTCT GCTGCGCTCG TCACCTTCTGG  
601 ACCAAAATCT GCAAACCGGT TCTGCACCAG GGTGAAGTTT GCACCAAACA  
651 GCGTAAAAAA GGTCCCCACG GTCTGGAAT CTTCCAGCGT TGCGACTGCG  
701 CTAAGGTCT GTCCTGCAA GTTGGAAAG ACGCTACCTA CTCCTCCAA  
751 GCTCGTCTGC ACGTTTGCCA GAAAATC

24 / 25

FIG. 24

1 ATGCAGCGTC TGGGTGCTAC CCTGCTGTGC CTGCTGCTGG CTGCTGCTGT  
51 TCCGACCGCT CCGGCTCCGG CTCCGACCGC TACCTCCGCT CCGTTAAAC  
101 CCGGTCCGGC TCTGTCTTAC CCGAGGAAG AAGCTACCTT GAACGAAATG  
151 TTCCGTGAAG TTGAAGAACT GATGAAGAC ACCAGCACA AACTGCGTTC  
201 CGCTGTTGAA GAAATGGAAG CTGAAGAAGC TGCTGCTAAA GCTTCTCCG  
251 AAGTTAACCT GGCTAACCTG CCGCCGTCTT ACCACAACGA AACCAACACC  
301 GACACCAAAG TTGGTAACAA CACCATCCAC GTTCACCGTG AAATCCACAA  
351 AATCACCAAC AACCAAGACG GTCAGATGGT TTTCTCCGAA ACCGTTATCA  
401 CCTCCGTTGG TGACGAAGAA GGTCTCGTT CCCACGAATG CATCATCGAC  
451 GAAGACTGG GTCCGTCCAT GTACTGCCAG TTCGCTTCCT TCCAGTACAC  
501 CTGCCAGCCG TGCCGTGGTC AGCGTATGCT GTGCACCCGT GACTCCGAAT  
551 GCTGCCGGTA CCAGCTGTGC GTTGGGGTC ACTGCACCAA AATGGCTACC  
601 CGTGGTTCCA ACGGTACCAT CTGCGACAAC CAGCGTACT GCCAGCCGGG  
651 TCTGTGCTGC GCTTTCAGC GTGGTCTGCT GTTCCCGGTT TGCACCCCGC  
701 TGCCGGTTGA AGGTGAAGT TGCCACGACC CGGCTTCCCG TCTGCTGGAC  
751 CTGATCACCT GGGAACTGA ACCGGACGGT GCTCTGGACC GTTGCCCGTG  
801 CGCTTCCGGT CTGCTGTGCC AGCCGCACTC CCACTCCCTG GTTTACGTTT  
851 GCAAAACCGAC CTTCGTTGGT TCCCGTGACC AGGACGGTGA AATCCTGCTG  
901 CCGCGTGAAG TTCCGGACGA ATACGAAGTT GGTTCCTTCA TGAAGAAGT  
951 TCGTCAGGAA CTGGAAGACC TGAACGTTT CCTGACCGAA GAAATGGCTC  
1001 TGGGTGAACC GGCTGCTGCT GCTGCTGCTC TGCTGGGTGG TGAAGAATC

25 / 25

## FIG. 25

1 ATGGTTGCTG CTGTTCTGCT GGGTCTGTCC TGGCTGTGCT CCCCCTGGG  
51 TGCTCTGGTT CTGGACTTCA ACAACATCCG TTCTTCCGCT GACCTGCACG  
101 GTGCTCGTAA AGGTTCCAG TGCCTGTCCG ACACCGACTG CAACACCCGT  
151 AAATTCTGCC TGCAGCCCGG TGACGAAAAA CCGTCTGCG CTACCTGCCG  
201 TGGTCTGCGT CGTCGTTGCC AGCGTGACGC TATGTGCTGC CCGGTACCC  
251 TGTGCGTTAA CGACGTTTGC ACCACCATGG AAGACGCTAC CCCGATCCTG  
301 GAACGTCAGC TGGACGAACA GGACGGTACC CACGCTGAAG GTACCACCGG  
351 TCACCCCGTT CAGGAAAACC AGCCGAAACG TAAACCGTCC ATCAAAAAAT  
401 CCCAGGGTCG TAAAGGTCAG GAAGGTGAAT CCTGCCCTGG TACCTTCGAC  
451 TGCGGTCCGG GTCTGTGCTG CGCTCGTCAC TTCTGGACCA AAATCTGCAA  
501 ACCGGTTCTG CTGGAAGGTC AGGTTTGCTC CCGTCGTGGT CACAAAGACA  
551 CCGCTCAGGC TCCGGAATC TTCCAGCGTT GCGACTGCGG TCCGGGTCTG  
601 CTGTGCCGTT CCCAGCTGAC CTCCAACCGT CAGCACGCTC GTCTGCGTGT  
651 TTGCCAGAAA ATCGAAAAAC TG

## SEQUENCE LISTING

&lt;110&gt; Amgen Inc.

&lt;120&gt; NOVEL DKR POLYPEPTIDES

&lt;130&gt; A-548

&lt;140&gt; 09/161,241

&lt;141&gt; 1998-09-25

&lt;160&gt; 78

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 1050

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 1

```
atgcagcggc tggggggtat ttgtctgtgt acactgctgg cggcgggcgggt ccccaactgct 60
cctgctcctt ccccgacggg cacttggact ccggcgggagc cggggcccagc tctcaactac 120
cctcaggagg aagctacgct caatgagatg ttctgagagg tggaggagct gatggaagac 180
actcagcaca aactgcgcag tgccgtggag gagatggagg cggagaagac agctgctaaa 240
acgtcctctg aggtgaacct ggcaagctta cctcccaact atcacaatga gaccagcacg 300
gagaccaggg tgggaaataa cacagtcctat gtgcaccagg aagttcacia gataaccaac 360
aaccagagtg gacaggtggg cttttctgag acagtcatta catctgtagg ggatgaagaa 420
ggcaagagga gccatgaatg tatcattgat gaagactgtg ggcccaccag gtactgccag 480
ttctccagct tcaagtacac ctgccagcca tgccggggacc agcagatgct atgcacccga 540
gacagtgagt gctgtggaga ccagctgtgt gcctgggggc actgcaccca aaaggccacc 600
aaagggtggc atgggaccat ctgtgacaac cagagggatt gccagcctgg cctgtgttgt 660
gccttccaaa gaggcctgct gttccccgtg tgcacacccc tgcccggtga gggagagctc 720
tgcatgacc ccaccagcca gctgctggat ctcatcacct gggaaactga gcctgaagga 780
gctttggacc gatgcccctg gcggcagtgcc ctccctatgcc agccacacag ccacagtctg 840
gtgtacatgt gcaagccagc cttcgtgggc agccatgacc acagtgaagg gagccagctg 900
cccagggagg ccccgatga gtacgaagat gttggcttca taggggaagt gcgccaggag 960
ctggaagacc tggagcggag cctagcccag gagatggcat ttgaggggccc tgcccctgtg 1020
gagtcactag gcggagagga ggagatttag                                     1050
```

&lt;210&gt; 2

&lt;211&gt; 1053

&lt;212&gt; DNA

&lt;213&gt; Human

&lt;400&gt; 2

```
atgcagcggc ttggggccac cctgctgtgc ctgctgctgg cggcgggcgggt cccacaggcc 60
cccgcgcccg ctccgacggc gacctcggct ccagtcaagc ccggcccggc tctcagctac 120
ccgaggagg aggccaccct caatgagatg ttccgagagg ttgaggaact gatggaggac 180
acgcagcaca aattgcgcag cgcggtggaa gagatggagg cagaagaagc tgctgctaaa 240
gcatcatcag aagtgaacct ggcaaaactta cctcccagct atcacaatga gaccaacaca 300
gacacgaagg ttggaaataa taccatccat gtgcaccgag aaattcacia gataaccaac 360
aaccagactg gacaaatggg cttttcagag acagttatca catctgtggg agacgaagaa 420
ggcagaagga gccacgagtg catcatcgac gaggactgtg ggcccagcat gtactgccag 480
tttgccagct tccagtacac ctgccagcca tgccgggggc agaggatgct ctgcacccgg 540
gacagtgagt gctgtggaga ccagctgtgt gtctgggggc actgcaccaa aatggccacc 600
```



aggggcagca atgggacat ctgtgacaac cagagggact gccagccggg gctgtgctgt 660  
gccttccaga gaggcctgt gttccctgtg tgcacacccc tgcccgtgga gggcgagctt 720  
tgccatgacc cggccagccg gcttctggac ctcatcacct gggagctaga gcctgatgga 780  
gccttggacc gatgcccttg tgccagtggc ctctctgcc agccccacag ccacagcctg 840  
gtgtatgtgt gcaagccgac ctctgtgggg agccgtgacc aagatgggga gatcctgtg 900  
cccagagagg tccccgatga gtatgaagtt ggcagcttca tggaggagggt gcgccaggag 960  
ctggaggacc tggagaggag cctgactgaa gagatggcgc tgggggagcc tgcggctgcc 1020  
gccgctgcac tgctgggagg ggaagagatt tag 1053

<210> 3  
<211> 801  
<212> DNA  
<213> Human

<400> 3  
atgatggctc tgggcgcagc gggagctacc cgggtctttg tcgcatgggt agcggcggct 60  
ctcggcggcc accctctgct gggagtgagc gccaccttga actcggttct caattccaac 120  
gctatcaaga acctgcccc accgctgggc ggcgctgcgg ggcacccagg ctctgcagtc 180  
agcgcgcgcg cgggaatcct gtaccgggc gggaataagt accagaccat tgacaactac 240  
cagccgtacc cgtgcgcaga ggacgaggag tgcggcactg atgagtactg cgctagtccc 300  
accgcgggag gggacgcggg cgtgcaaacc tgtctcgct gcaggaagcg ccgaaaacgc 360  
tgcatgcgtc acgctatgtg ctgccccggg aattactgca aaaatggaat atgtgtgtct 420  
tctgatcaaa atcatttccg aggagaaatt gaggaaccca tcaactgaaag ctttggtaat 480  
gatcatagca ccttggatgg gtattccaga agaaccacct tgtcttcaaa aatgtatcac 540  
accaaaggac aagaaggttc tgtttgtctc cggtcacag actgtgcctc aggtattgtg 600  
tgtgctagac acttctggtc caagatctgt aaacctgtcc tgaaagaagg tcaagtgtgt 660  
accaagcata ggagaaaagg ctctcatgga ctagaaatat tccagcgttg ttactgtgga 720  
gaaggtctgt cttgccggat acagaaagat caccatcaag ccagtaattc ttctaggctt 780  
cacactgtc agagacacta a 801

<210> 4  
<211> 780  
<212> DNA  
<213> Mouse

<400> 4  
atggccgcgc tgatgcgggt caaggattca tcccgtgcc ttctctact ggccgcgggtg 60  
ctgatgggtg agagctcaca gctaggcagc tcgcccggcca aactcaactc catcaagtcc 120  
tctctaggag gggagactcc tgctcagtca gccaaaccgat ctgcaggcat gaaccaagga 180  
ctggctttcg gcggcagtaa gaagggcaaa agcctggggc aggcctaccc ttgcagcagt 240  
gataaggaat gtgaagtgg aagatactgc cacagtcccc accaaggatc atcagcctgc 300  
atgctctgta ggaggaaaaa gaaacgatgc cacagagatg ggatgtgttg ccctggtacc 360  
cgctgcaata atggaatctg catcccagtc actgagagca tcctcacccc acatatccca 420  
gctctggatg gcacccggca tagagatcgc aaccatggtc actattccaa ccatgacctg 480  
ggatggcaga atctaggaag gccacactcc aagatgcctc atataaaagg acatgaagga 540  
gacccatgcc tacggtcatc agactgcatt gatgggtttt gttgtgctcg ccacttctgg 600  
acaaaaatct gcaaaccagt gctccatcag ggggaagtct gtaccaaaca acgcaagaag 660  
ggttcgcacg ggctggagat tttccagagg tgtgactgtg caaagggcct gtccctgcaa 720  
gtgtggaaag atgccaccta ctcttccaaa gccagactcc atgtatgcca gaagatctga 780

<210> 5  
<211> 780  
<212> DNA  
<213> Human

<400> 5  
atggccgcgt tgatgcggag caaggattcg tctgtctgcc tgctcctact ggccgcgggtg 60

ctgatggtgg agagctcaca gatcggcagt tcgcgggcca aactcaactc catcaagtcc 120  
tctctgggcg gggagacgcc tggtcaggcc gccaatcgat ctgcgggcat gtaccaagga 180  
ctggcattcg gcggcagtaa gaagggcaaa aacctggggc aggcctaccc ttgtagcagt 240  
gataaggagt gtgaagtggg gaggtattgc cacagtcccc accaaggatc atcggcctgc 300  
atggtgtgtc ggagaaaaaa gaagcgctgc caccgagatg gcatgtgctg ccccgatcc 360  
cgctgcaata atggcatctg tatcccagtt actgaaagca tcttaacccc tcacatcccc 420  
gctctggatg gtactcggca cagagatcga aaccacggtc attactcaaa ccatgacttg 480  
ggatggcaga atctaggaag accacacact aagatgtcac atataaaagg gcatgaagga 540  
gacccctgcc tacgatcatc agactgcatt gaagggtttt gctgtgctcg tcatttctgg 600  
acaaaaatct gcaaacacagt gctccatcag ggggaagtct gtaccaaaca acgcaagaag 660  
ggttctcatg ggctggaaat tttccagcgt tgcgactgtg cgaagggcct gtcttgcaaa 720  
gtatggaaag atgccaccta ctctccaaa gccagactcc atgtgtgtca gaaaatttga 780

<210> 6  
<211> 624  
<212> DNA  
<213> Human

<400> 6  
atggccgcgt tgatcgggag caaggattcg tctgtctgcc tgctcctact ggccgcggtg 60  
ctgatggtgg agagctcaca gatcggcagt tcgcgggcca aactcaactc catcaagtcc 120  
tctctgggcg gggagacgcc tggtcaggcc gccaatcgat ctgcgggcat gtaccaagga 180  
ctggcattcg gcggcagtaa gaagggcaaa aacctggggc aggcctaccc ttgtagcagt 240  
gataaggagt gtgaagtggg gaggtattgc cacagtcccc accaaggatc atcggcctgc 300  
atggtgtgtc ggagaaaaaa gaagcgctgc caccgagatg gcatgtgctg ccccgatcc 360  
cgctgcaata atgggcatga aggagacccc tgcctacgat catcagactg cattgaaggg 420  
ttttgctgtg ctctgcatct ctggacaaaa atctgcaaac cagtgtctca tcagggggaa 480  
gtctgtacca aacaacgcaa gaagggttct catgggctgg aaattttcca gcgttgcgac 540  
tgtgcgaagg gcctgtcttg caaagtatgg aaagatgcca cctactcctc caaagccaga 600  
ctccatgtgt gtcagaaaat ttga 624

<210> 7  
<211> 675  
<212> DNA  
<213> Human

<400> 7  
atggtggcgg ccgtcctgct ggggctgagc tggctctgct ctccccctggg agctctggtc 60  
ctggacttca acaacatcag gagctctgct gacctgcatg gggcccggaa gggctcacag 120  
tgctgtctg acacggactg caataccaga aagttctgcc tccagccccg cgatgagaag 180  
ccgttctgtg ctacatgtcg tgggttgctg aggaggtgcc agcgagatgc catgtgctgc 240  
cctgggacac tctgtgtgaa cgatgtttgt actacgatgg aagatgcaac cccaatatta 300  
gaaaggcagc ttgatgagca agatggcaca catgcagaag gaacaactgg gcacccagtc 360  
caggaaaacc aacccaaaag gaagccaagt attaagaaat cacaaggcag gaagggacaa 420  
gagggagaaa gttgtctgag aacttttgac tgtggccctg gactttgctg tgctcgtcat 480  
ttttggacga aaatttgtaa gccagtcctt ttggaggggac aggtctgctc cagaagaggg 540  
cataaagaca ctgctcaagc tccagaaatc ttccagcgtt gcgactgtgg ccctggacta 600  
ctgtgtcgaa gccaatgac cagcaatcgg cagcatgctc gattaagagt atgccaaaaa 660  
atagaaaagc tataa 675

<210> 8  
<211> 349  
<212> PRT  
<213> Mouse

&lt;400&gt; 8

Met Gln Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala  
1 5 10 15

Val Pro Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala  
20 25 30

Glu Pro Gly Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn  
35 40 45

Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys  
50 55 60

Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys  
65 70 75 80

Thr Ser Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn  
85 90 95

Glu Thr Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His  
100 105 110

Gln Glu Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe  
115 120 125

Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser  
130 135 140

His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln  
145 150 155 160

Phe Ser Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met  
165 170 175

Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp  
180 185 190

Gly His Cys Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys  
195 200 205

Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg  
210 215 220

Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu  
225 230 235 240

Cys His Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu  
245 250 255

Glu Pro Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu  
260 265 270

Cys Gln Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe  
275 280 285

Val Gly Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala  
290 295 300

Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu  
305 310 315 320

Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly  
325 330 335

Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile  
340 345

<210> 9  
<211> 350  
<212> PRT  
<213> Human

<400> 9  
Met Gln Arg Leu Gly Ala Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala  
1 5 10 15

Val Pro Thr Ala Pro Ala Pro Ala Pro Thr Ala Thr Ser Ala Pro Val  
20 25 30

Lys Pro Gly Pro Ala Leu Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn  
35 40 45

Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys  
50 55 60

Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Lys  
65 70 75 80

Ala Ser Ser Glu Val Asn Leu Ala Asn Leu Pro Pro Ser Tyr His Asn  
85 90 95

Glu Thr Asn Thr Asp Thr Lys Val Gly Asn Asn Thr Ile His Val His  
100 105 110

Arg Glu Ile His Lys Ile Thr Asn Asn Gln Thr Gly Gln Met Val Phe  
115 120 125

Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Arg Arg Ser  
130 135 140

His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln  
145 150 155 160

Phe Ala Ser Phe Gln Tyr Thr Cys Gln Pro Cys Arg Gly Gln Arg Met  
165 170 175

Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Val Trp  
180 185 190

Gly His Cys Thr Lys Met Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys  
195 200 205

Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg  
 210 215 220  
 Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu  
 225 230 235 240  
 Cys His Asp Pro Ala Ser Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu  
 245 250 255  
 Glu Pro Asp Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu  
 260 265 270  
 Cys Gln Pro His Ser His Ser Leu Val Tyr Val Cys Lys Pro Thr Phe  
 275 280 285  
 Val Gly Ser Arg Asp Gln Asp Gly Glu Ile Leu Leu Pro Arg Glu Val  
 290 295 300  
 Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu  
 305 310 315 320  
 Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Gly Glu  
 325 330 335  
 Pro Ala Ala Ala Ala Ala Ala Leu Leu Gly Gly Glu Glu Ile  
 340 345 350

<210> 10  
 <211> 266  
 <212> PRT  
 <213> Human

<400> 10  
 Met Met Ala Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met  
 1 5 10 15  
 Val Ala Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr  
 20 25 30  
 Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro  
 35 40 45  
 Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro  
 50 55 60  
 Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr  
 65 70 75 80  
 Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr  
 85 90 95  
 Cys Ala Ser Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys Leu  
 100 105 110  
 Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys Cys  
 115 120 125

Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln Asn  
 130 135 140  
 His Phe Arg Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly Asn  
 145 150 155 160  
 Asp His Ser Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser Ser  
 165 170 175  
 Lys Met Tyr His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg Ser  
 180 185 190  
 Ser Asp Cys Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser Lys  
 195 200 205  
 Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His Arg  
 210 215 220  
 Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys Gly  
 225 230 235 240  
 Glu Gly Leu Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser Asn  
 245 250 255  
 Ser Ser Arg Leu His Thr Cys Gln Arg His  
 260 265

<210> 11  
 <211> 259  
 <212> PRT  
 <213> Mouse

<400> 11  
 Met Ala Ala Leu Met Arg Val Lys Asp Ser Ser Arg Cys Leu Leu Leu  
 1 5 10 15  
 Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Leu Gly Ser Ser Arg  
 20 25 30  
 Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly Glu Thr Pro Ala  
 35 40 45  
 Gln Ser Ala Asn Arg Ser Ala Gly Met Asn Gln Gly Leu Ala Phe Gly  
 50 55 60  
 Gly Ser Lys Lys Gly Lys Ser Leu Gly Gln Ala Tyr Pro Cys Ser Ser  
 65 70 75 80  
 Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser Pro His Gln Gly  
 85 90 95  
 Ser Ser Ala Cys Met Leu Cys Arg Arg Lys Lys Lys Arg Cys His Arg  
 100 105 110

Asp Gly Met Cys Cys Pro Gly Thr Arg Cys Asn Asn Gly Ile Cys Ile  
           115                                  120                                  125  
 Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly  
           130                                  135                                  140  
 Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu  
 145                                  150                                  155                                  160  
 Gly Trp Gln Asn Leu Gly Arg Pro His Ser Lys Met Pro His Ile Lys  
                                   165                                  170                                  175  
 Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Asp Gly  
                                   180                                  185                                  190  
 Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu  
                                   195                                  200                                  205  
 His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly  
                                   210                                  215                                  220  
 Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys  
 225                                  230                                  235                                  240  
 Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys  
                                   245                                  250                                  255  
 Gln Lys Ile

<210> 12  
 <211> 259  
 <212> PRT  
 <213> Human

<400> 12  
 Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys Cys Leu Leu Leu  
   1                                  5                                  10                                  15  
 Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile Gly Ser Ser Arg  
                                   20                                  25                                  30  
 Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly Glu Thr Pro Gly  
                                   35                                  40                                  45  
 Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly Leu Ala Phe Gly  
                                   50                                  55                                  60  
 Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr Pro Cys Ser Ser  
                                   65                                  70                                  75                                  80  
 Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser Pro His Gln Gly  
                                   85                                  90                                  95  
 Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg  
                                   100                                  105                                  110

Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile  
           115                                  120                                  125  
 Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly  
           130                                  135                                  140  
 Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu  
   145                                  150                                  155                                  160  
 Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys  
                                   165                                  170                                  175  
 Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly  
                                   180                                  185                                  190  
 Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu  
                                   195                                  200                                  205  
 His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly  
                                   210                                  215                                  220  
 Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys  
   225                                  230                                  235                                  240  
 Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys  
                                   245                                  250                                  255

Gln Lys Ile

<210> 13  
 <211> 207  
 <212> PRT  
 <213> Human

<400> 13  
 Met Ala Ala Leu Met Arg Ser Lys Asp Ser Ser Cys Cys Leu Leu Leu  
   1                                  5                                  10                                  15  
 Leu Ala Ala Val Leu Met Val Glu Ser Ser Gln Ile Gly Ser Ser Arg  
                                   20                                  25                                  30  
 Ala Lys Leu Asn Ser Ile Lys Ser Ser Leu Gly Gly Glu Thr Pro Gly  
                                   35                                  40                                  45  
 Gln Ala Ala Asn Arg Ser Ala Gly Met Tyr Gln Gly Leu Ala Phe Gly  
                                   50                                  55                                  60  
 Gly Ser Lys Lys Gly Lys Asn Leu Gly Gln Ala Tyr Pro Cys Ser Ser  
   65                                  70                                  75                                  80  
 Asp Lys Glu Cys Glu Val Gly Arg Tyr Cys His Ser Pro His Gln Gly  
                                   85                                  90                                  95



Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg  
                   100                                  105                                  110  
 Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly His Glu Gly  
                   115                                  120                                  125  
 Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly Phe Cys Cys Ala  
                   130                                  135                                  140  
 Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu His Gln Gly Glu  
                   145                                  150                                  155                                  160  
 Val Cys Thr Lys Lys Arg Lys Lys Gly Ser His Gly Leu Glu Ile Phe  
                   165                                  170                                  175  
 Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys Val Trp Lys Asp  
                   180                                  185                                  190  
 Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys Gln Lys Ile  
                   195                                  200                                  205

<210> 14  
 <211> 224  
 <212> PRT  
 <213> Human

<400> 14  
 Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro Leu  
   1                                  5                                  10                                  15  
 Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp Leu  
                   20                                  25                                  30  
 His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys Asn  
                   35                                  40                                  45  
 Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys Ala  
                   50                                  55                                  60  
 Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys Cys  
                   65                                  70                                  75                                  80  
 Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp Ala  
                   85                                  90                                  95  
 Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His Ala  
                   100                                  105                                  110  
 Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg Lys  
                   115                                  120                                  125  
 Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu Ser  
                   130                                  135                                  140  
 Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg His  
                   145                                  150                                  155                                  160

Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val Cys  
165 170 175  
Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe Gln  
180 185 190  
Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser  
195 200 205  
Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu  
210 215 220

<210> 15  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 15  
ggaaggaaaa aagcgggccgc aacannnnnnn nnn

33

<210> 16  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide adapter

<400> 16  
tcgacccacg cgtccg

16

<210> 17  
<211> 12  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide adapter

<400> 17  
gggtgcgag gc

12

<210> 18  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 18

actagctcca gtgatctc

18

<210> 19

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 19

cgtcattgtt ctcgttcc

18

<210> 20

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 20

ccagctgctc tgtggcagcc cag

23

<210> 21

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 21

cccagtcacg acgttgtaaa acgacggcc

29

<210> 22

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 22

aacatgcagc ggctcggggg

20

<210> 23  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 23  
ggtgacacta tagaagagct atgacgtcgc

30

<210> 24  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 24  
gtgctgagtg tcttccatca gc

22

<210> 25  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probes

<400> 25  
gagatgcagc ggcttggggc caccc

25

<210> 26  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probes

<400> 26  
gcctggtcag cccacgccta aag

23

<210> 27  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probes

<400> 27  
cctgctgctg gcggcgccgg tccccacggc 30

<210> 28  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probes

<400> 28  
gcctggtcag cccacgccta aag 23

<210> 29  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probes

<400> 29  
cccggaccct gactctgcag ccg 23

<210> 30  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probes

<400> 30  
gaggaaaaat aggcagtgcg gcacc 25

<210> 31  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide primers

<400> 31  
gccacagtcc ccaccaagga tcac 25

<210> 32  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primers

<400> 32

gatgacctt ggtggggact gtggc

25

<210> 33

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primers

<400> 33

ctgcaaacca gtgctccatc aggg

24

<210> 34

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primers

<400> 34

ccctgatgga gcactggttt gcag

24

<210> 35

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primer

<400> 35

gctataccaa gcatacaatc

20

<210> 36

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 36

gggttgaggg aacacaatct gcaag

25

<210> 37

<211> 28

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 37  
gtctgcaatt gatgatgttc ctcaatgg

28

<210> 38  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 38  
ccagggccac agtcgcaacg ctgg

24

<210> 39  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 39  
ctccctcttg tcccttcctg ccttg

25

<210> 40  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 40  
caaggcagga agggacaaga gggag

25

<210> 41  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 41  
ccagcggttg gactgtggcc ctgg

24

<210> 42  
<211> 44  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide primer/adapter

<400> 42  
gactagtcttct agatcgcgag cggccgcctt tttttttttt tttt

44

<210> 43  
<211> 6  
<212> PRT  
<213> Human

<400> 43  
Met His Pro Leu Leu Gly  
1 5

<210> 44  
<211> 5  
<212> PRT  
<213> Human

<400> 44  
Thr Cys Gln Arg His  
1 5

<210> 45  
<211> 59  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 45  
gttctctca tatgcatcca ttattaggcg taagtgccac cttgaactcg gttctcaat 59

<210> 46  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 46  
tacgcactgg atccttagtg tctctgacaa gtgtgaag

38



<210> 47  
<211> 6  
<212> PRT  
<213> Human

<400> 47  
Met Ser Gln Ile Gly Ser  
1 5

<210> 48  
<211> 5  
<212> PRT  
<213> Human

<400> 48  
Val Cys Gln Lys Ile  
1 5

<210> 49  
<211> 56  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 49  
gttctctca tatgtctcaa attgtagtt ctctgtgcaa actcaactcc atcaag 56

<210> 50  
<211> 39  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 50  
tacgcactgg atccttaaatt tttctgacac acatggagt 39

<210> 51  
<211> 6  
<212> PRT  
<213> Mouse

<400> 51  
Met Ser Gln Leu Gly Ser  
1 5

<210> 52  
<211> 5  
<212> PRT

<213> Mouse

<400> 52

Val Cys Gln Lys Ile  
1 5

<210> 53

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 53

gttctctca tatgtctcaa ttaggtagct ctcgtgctaa actcaactcc atcaagtcc 59

<210> 54

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 54

tacgcactgg atccttagat cttctggcat acatggagt 39

<210> 55

<211> 6

<212> PRT

<213> Human

<400> 55

Met Pro Ala Pro Thr Ala  
1 5

<210> 56

<211> 5

<212> PRT

<213> Human

<400> 56

Gly Gly Glu Glu Ile  
1 5

<210> 57

<211> 54

<212> DNA

<213> Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide probe

&lt;400&gt; 57

gttctctctca tatgcctgct ccaactgcaa cttcggctcc agtcaagccc ggcc

54

&lt;210&gt; 58

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide probe

&lt;400&gt; 58

tagcactcc gcggttaaatt ctcttcccct cccagca

37

&lt;210&gt; 59

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Human

&lt;400&gt; 59

Met Lys Pro Gly Pro Ala

1

5

&lt;210&gt; 60

&lt;211&gt; 5

&lt;212&gt; PRT

&lt;213&gt; Human

&lt;400&gt; 60

Gly Gly Glu Glu Ile

1

5

&lt;210&gt; 61

&lt;211&gt; 54

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide probe

&lt;400&gt; 61

gttctctctca tatgaaacca ggtccagcct taagctaccc gcaggaggag gccca

54

&lt;210&gt; 62

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide probe

&lt;400&gt; 62

tacgcactcc gcggttaaatt ctcttcccct cccagca

37

&lt;210&gt; 63

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Human

&lt;400&gt; 63

Met Gln Glu Glu Ala Thr

1 5

&lt;210&gt; 64

&lt;211&gt; 5

&lt;212&gt; PRT

&lt;213&gt; Human

&lt;400&gt; 64

Gly Gly Glu Glu Ile

1 5

&lt;210&gt; 65

&lt;211&gt; 53

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide probe

&lt;400&gt; 65

gttctctca tatgcaagaa gaagctactc tgaatgagat gttccgcgag gtt

53

&lt;210&gt; 66

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
Oligonucleotide probe

&lt;400&gt; 66

tacgcactcc gcggttaaatt ctcttcccct cccagca

37

&lt;210&gt; 67

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Mouse

<400> 67  
Met Glu Pro Gly Pro Ala  
1 5

<210> 68  
<211> 5  
<212> PRT  
<213> Mouse

<400> 68  
Gly Glu Glu Glu Ile  
1 5

<210> 69  
<211> 54  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 69  
gttctcctca tatggaacca ggtccagctt taaactaccc tcaggaggaa gcta 54

<210> 70  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 70  
tacgcactcc gcggttaaatt ctctcctctt ccgccta 37

<210> 71  
<211> 6  
<212> PRT  
<213> Human

<400> 71  
Met Leu Val Leu Asp Phe  
1 5

<210> 72  
<211> 5  
<212> PRT  
<213> Human

<400> 72  
Lys Ile Glu Lys Leu  
1 5

<210> 73  
<211> 47  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 73  
gttctcctca tatgtagtt ttggatttca acaacatcag gagctct 47

<210> 74  
<211> 49  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide probe

<400> 74  
tacgcactgg atccttacag ttttcttatt ttttggcata ctcttaatc 49

<210> 75  
<211> 798  
<212> DNA  
<213> Human

<400> 75  
atgatggctc tgggtgctgc tgggtgctacc cgtggttttcg ttgctatggg tgcgtgctgct 60  
ctgggtgggc acccgctgct ggggtgtttcc gctaccctga actccgttct gaactccaac 120  
gctatcaaaa acctgccgcc gccgctgggt ggtgctgctg gtcaccggg ttccgctggt 180  
tccgctgctc cgggtatcct gtaccgggt ggtaacaaat accagaccat cgacaactac 240  
cagccgtacc cgtgcgctga agacgaagaa tgcgggtaccg acgaatactg cgcttccccg 300  
acccgtgggtg gtgacgctgg tgttcagatc tgccctggctt gccgtaaacg tcgtaaacgt 360  
tgcatgcgtc acgctatgtg ctgcccgggt aactactgca aaaacgggtat ctgcgtttcc 420  
tccgaccaga accacttccg tggtgaaatc gaagaaacca tcaccgaatc cttcggtaac 480  
gaccactcca ccctggacgg ttactccgt cgtaccacc tgcctccaa aatgtaccac 540  
accaaaggctc aggaagggtc cgtttgctg cgttccctccg actgcgcttc cggctctgtgc 600  
tgcgctcgtc acttctgggtc caaaatctgc aaaccgggtc tgaaagaagg tcagggtttgc 660  
accaaacacc gtcgtaaaagg ttcccacgggt ctggaaatct tccagcgttg ctactgcgggt 720  
gaaggctctgt cctgccgtat ccagaaagac caccaccagg cttccaactc ctcccgtctg 780  
cacacctgcc agcgtcac 798

<210> 76  
<211> 777  
<212> DNA  
<213> Human

<400> 76  
atggctgctc tgatgcgttc caaagactcc tctgctgccc tgcgtgctgct ggctgctggt 60  
ctgatgggtg aatcctccca gatcgggtcc tcccgctgcta aactgaactc catcaaatcc 120  
tccctgggtg gtgaaacccc gggtcagggt gctaaccggt ccgctggtat gtaccagggt 180  
ctggcttttcg gtggttccaa aaaaggtaaa aacctgggtc aggtttaccg gtgctcctcc 240

```
gacaaagaat gCGaagttgg tcgttactgc cactccccgc accagggttc ctccggttgc 300
atggttttgcc gtcgtaaaaa aaaacgttgc caccgtgacg gtatgtgctg cccgtccacc 360
cgttgcaaca acggtatctg catcccggtt accgaatcca tcttgacccc gcacatcccc 420
gctctggacg gtaccgctca ccgtgaccgt aaccacggtc actactccaa ccacgacctg 480
ggttggcaga acctgggtcg tccgcacacc aaaatgtccc acatcaaagg tcacgaaggt 540
gacccggtgc tgcgttcctc cgactgcacg gaaggtttct gctgcgctcg tcacttctgg 600
accaaaatct gcaaaccggt tctgcaccag ggtgaagttt gcaccaaaca gcgtaaaaaa 660
ggttcccacg gtctggaaat cttccagcgt tgcgactgcg ctaaagggtc gtccctgcaa 720
gtttggaaag acgctaccta ctctccaaa gctcgtctgc acgtttgcca gaaaatc 777
```

&lt;210&gt; 77

&lt;211&gt; 1050

&lt;212&gt; DNA

&lt;213&gt; Human

&lt;400&gt; 77

```
atgcagcgtc tgggtgctac cctgctgtgc ctgctgctgg ctgctgctgt tccgaccgct 60
ccggtcccg ctcgaccgc tacctccgct ccggttaaac cgggtccggc tctgtcctac 120
ccgcaggaag aagctaccct gaacgaaatg ttccgtgaag ttgaagaact gatggaagac 180
accagcaca aactgcgttc cgctgttgaa gaaatggaag ctgaagaagc tgctgctaaa 240
gcttccctcg aagttaacct ggctaacctg ccgcgcctct accacaacga aaccaacacc 300
gacacaaaag ttggtataca caccatccac gttcacctg aaatccacaa aatcaacaa 360
aaccagaccg gtcagatggt tttctccgaa accgttatca ctccggttg tgacgaagaa 420
ggctcgtcgt cccacgaatg catcatcgac gaagactgcg gtccgtccat gtactgccag 480
ttcgttccct tccagtacac ctgccagccg tgcctgggtc agcgtatgct gtgcaccctg 540
gactccgaat gctgcgggtg ccagctgtgc gtttggggtc actgcaccaa aatgggtacc 600
cgtggttcca acggtaccat ctgcgacaac cagcgtgact gccagccggg tctgtgctgc 660
gcttccagc gtggtctgct gttcccggtt tgcacccgc tgcgggttg aggtgaactg 720
tgccacgacc cggcttcccg tctgctggac ctgatcacct gggaaactga accggacggt 780
gctctggacc gttgcccgtg cgcttccggt ctgctgtgcc agccgcactc cactccctg 840
gtttacgttt gcaaaccgac cttcgttggt tcccgtgacc aggacggtga aatcctgctg 900
ccgctgaag ttccggacga atacgaagtt ggttccttca tggagaagt tcgtcaggaa 960
ctggaagacc tggaaacgtt cctgaccgaa gaaatggctc tgggtgaacc ggctgctgct 1020
gctgctgctc tgctgggtgg tgaagaaatc 1050
```

&lt;210&gt; 78

&lt;211&gt; 672

&lt;212&gt; DNA

&lt;213&gt; Human

&lt;400&gt; 78

```
atggttgctg ctgttctgct gggctctgtc tggctgtgct ccccgctggg tgctctggtt 60
ctggacttca acaacatccg ttccctccgct gacctgcacg gtgctcgtaa aggttcccag 120
tgctgtcctg acaccgactg caacaccctg aaattctgcc tgcagcccg tgacgaaaaa 180
ccgttctgct ctacctgccc tgggtctgct cgctggtgcc agcgtgacgc tatgtgctgc 240
ccgggtaccc tgtgcgttaa cgacgtttgc accaccatgg aagacgctac cccgatcctg 300
gaacgtcagc tggacgaaca ggacggtacc cagcgtgaag gtaccaccgg tcaccgggtt 360
caggaaaaac agccgaaacg taaaccgtcc atcaaaaaat cccagggctg taaaggctag 420
gaaggtgaat cctgcctgctg taccttccgac tgcgggtccg gtctgtgctg cgctcgtcac 480
ttctggacca aaatctgcaa accggttctg ctggaaggct aggtttgctc ccgtcgtggt 540
cacaagagca ccgctcaggc tccggaaatc ttccagcgtt gcgactgctg tccgggtctg 600
ctgtgccgtt cccagctgac ctccaaccgt cagcacgctc gtctgcgtgt ttgccagaaa 660
atcgaaaaac tg 672
```

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/12, C07K 14/475</b>		<b>A3</b>	(11) International Publication Number: <b>WO 00/18914</b>
			(43) International Publication Date: 6 April 2000 (06.04.00)
(21) International Application Number: <b>PCT/US99/21647</b>		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 17 September 1999 (17.09.99)		<b>Published</b> <i>With international search report.</i>	
(30) Priority Data: 09/161,241 25 September 1998 (25.09.98) US		(88) Date of publication of the international search report: 31 August 2000 (31.08.00)	
(71) Applicant: AMGEN INC. [US/US]; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).			
(72) Inventors: BASS, Michael, Brian; 2208 Rustic Park Court, Thousand Oaks, CA 91362 (US). SULLIVAN, John, Kevin; 1085 Rotella Street, Newbury Park, CA 91320 (US). THEILL, Lars, Eyde; 1874 Calle Borrego, Thousand Oaks, CA 91360 (US). WANG, Daguang; 1800 Via Petirrojo #K, Thousand Oaks, CA 91320 (US).			
(74) Agents: ODRE, Steven, M. et al.; Amgen, Inc., One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).			
(54) Title: NOVEL <i>DKR</i> POLYPEPTIDES			
1 MQRLGATLLC LLLAAAVPTA PAPAPTATSA PVKPGPALSYPQEEATLNEM 51 FREVEELMED TQHKLRSAVE EMEAEAAAAK ASSEVNLANL PPSYHNETNT 101 DTKVGNNTIH VHREIHKITN NQTGQMVFSE TVITSVGDEE GRRSHECIID 151 EDCGPSMYCQ FASFQYTCQP CRGQRM LCTR DSECCGDQLC VWGHCTKMAT 201 RGSNGTICDN QRDCQPGLCC AFQRG L LFPV CTPLPVEGEL CHDPASRLLD 251 LITWELEPDG ALDRCPCASG LLCQPHSHSL VYVCKPTFVG SRDQDGEILL 301 PREVPDEYEV GSFMEEVROE LEDLERSLTE EMALGEPAAA AAALLGGEET 351 *			
(57) Abstract			
Disclosed are nucleic acid molecules encoding novel <i>DKR</i> polypeptides. Also disclosed are methods of preparing the nucleic acid molecules and polypeptides, and methods of using these molecules.			



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/21647

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/475

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 98 27932 A (HUMAN GENOME SCIENCES INC ; RUBEN STEVEN M (US); SOPPET DANIEL R (U) 2 July 1998 (1998-07-02) page 3, line 22 - page 4, line 5 page 8, line 6 - line 11 page 8, line 19 - page 9, line 16 claims 1-20; examples 1-4 --- -/-</p>	1-20

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

15 May 2000

Date of mailing of the international search report

31. 05. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

van Klompenburg, W

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/21647

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GLINKA ET AL: "Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction" NATURE,GB,MACMILLAN JOURNALS LTD. LONDON, vol. 391, no. 6665, 22 January 1998 (1998-01-22), pages 357-362, XP002096088 ISSN: 0028-0836 page 357, column 1 -column 2 page 360, column 1 -column 2, paragraph 1 page 361, column 2 -page 362 figures 1,5 -& STRAUSBERG ET AL.: "National cancer institute, cancer genome anatomy project (CGAP)" EMBL DATABASE ACC NO: AA207078, 31 January 1997 (1997-01-31), XP002137654 ---	1-20
X	MARRA ET AL.: "The WashU-HMMI Mouse EST project" EMBL DATABASE ACC NO:AA265561, 21 March 1997 (1997-03-21), XP002137655 abstract ---	1-20
X	HILLIER ET AL.: "The WashU-Merck EST project" EMBL DATABASE ACC NO: W55979, 6 June 1996 (1996-06-06), XP002137656 abstract ---	1-20
X	STRAUSBERG ET AL.: "National cancer institute, cancer genome anatomy project (CGAP)" EMBL DATABASE ACC NO: AA565546, 11 September 1997 (1997-09-11), XP002137657 abstract ---	1-20
A	WO 97 48275 A (UNIV CALIFORNIA) 24 December 1997 (1997-12-24) page 4, line 21 - line 33 page 6, line 6 - line 9 page 12, line 25 -page 13, line 29 page 14, line 21 - line 22; claims 1-15 ---	1-20
P,X	WO 98 46755 A (MCCARTHY SEAN A ;MILLENNIUM BIOTHERAPEUTICS INC (US)) 22 October 1998 (1998-10-22) page 1, line 1 - line 34 SEQ ID NOs: 1,2,16 & 17 page 12, line 12 -page 13, line 6; claims 1-22; figures 6,7 ---	1-20
	---	
	-/--	

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/21647

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>WO 99 14328 A (CHEN JIAN ;GENENTECH INC (US); PENNICA DIANE (US); YUAN JEAN (US);) 25 March 1999 (1999-03-25) page 1 page 26, line 28 -page 27, line 2 page 48, line 4 - line 14 page 58, line 36 -page 59, line 2 page 144, line 10 - line 37 page 185, line 9 - line 22 SEQ ID NOs: 235 &amp; 236 claims 1-18; figures 83,84 ---</p>	1-20
P,X	<p>TATE ET AL.: "Homo sapiens hddk-4 mRNA, complete cds." EMBL DATABASE ACC NO : AB017788, 29 September 1998 (1998-09-29), pages 239-242, XP002137658 abstract -----</p>	1-20

# INTERNATIONAL SEARCH REPORT

national application No.  
PCT/US 99/21647

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/21647

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-20 all partially

An isolated nucleic acid molecule encoding a polypeptide, comprising any of SEQ ID NOs: 1,2 and 77 and a nucleic acid molecule that hybridizes to the above under stringent conditions. A process for producing a polypeptide, preferably SEQ ID NOs: 8 and 9, comprising expressing a polypeptide encoded by the above mentioned nucleic acid in a suitable host and isolating the polypeptide.

An isolated nucleic acid molecule which is the complement of any of the above mentioned nucleic acids.

A nucleic acid molecule encoding any of the polypeptides of: SEQ ID NOs 8 and 9, or a polypeptide that has 1-100 amino acid substitutions and/or deletions as compared to said polypeptides, or a fragment of SEQ ID NOs: 8 and 9. A vector comprising any of the above mentioned nucleic acid molecules. A polypeptide of SEQ ID NOs: 8 or 9, or said polypeptide that does not possess an endogenous signal peptide, or a polypeptide that is at least 85% identical to said polypeptide or a fragment of said polypeptide.

2. Claims: 1-20 all partially

Idem for SEQ ID NOs 3, 10 and 75

3. Claims: 1-20 all partially

Idem for SEQ ID NOs: 4,5,6,11,12,13 and 76

4. Claims: 1-20 all partially

Idem for SEQ ID NOs: 7, 14 and 78

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/US 99/21647

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9827932 A	02-07-1998	AU 5613498 A EP 0954575 A	17-07-1998 10-11-1999
WO 9748275 A	24-12-1997	AU 3576597 A CA 2258789 A EP 0973391 A	07-01-1998 24-12-1997 26-01-2000
WO 9846755 A	22-10-1998	AU 7137398 A EP 0975755 A	11-11-1998 02-02-2000
WO 9914328 A	25-03-1999	AU 9317898 A AU 9312198 A AU 9484398 A WO 9914327 A WO 9914234 A AU 9395998 A WO 9914241 A AU 9317498 A AU 1126099 A AU 1288399 A WO 9921998 A WO 9921999 A AU 1532499 A WO 9927098 A AU 1703399 A WO 9927100 A	05-04-1999 05-04-1999 05-04-1999 25-03-1999 25-03-1999 05-04-1999 25-03-1999 05-04-1999 17-05-1999 17-05-1999 06-05-1999 06-05-1999 15-06-1999 03-06-1999 15-06-1999 03-06-1999

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**